

Chi-boundedness of graphs containing no cycles with k chords

Joonkyung Lee*

Shoham Letzter[†]

Alexey Pokrovskiy[‡]

Abstract

We prove that the family of graphs containing no cycle with exactly k -chords is χ -bounded, for k large enough or of form $\ell(\ell - 2)$ with $\ell \geq 3$ an integer. This verifies (up to a finite number of values k) a conjecture of Aboulker and Bousquet (2015).

1 Introduction

The *clique number* of a graph G , denoted $\omega(G)$, is the size of its largest clique in G . The *chromatic number* of G , denoted $\chi(G)$, is the minimum number of colours in a *proper vertex-colouring* of G , which is a colouring of the vertices where adjacent vertices have distinct colours. It is easy to see that $\chi(G) \geq \omega(G)$ for every graph G , but the converse is far from the truth. Indeed, the chromatic number cannot be upper-bounded by a function of the clique number. This can be seen, for example, through a construction due to Mycielski [12] that provides a family of triangle-free graphs whose chromatic number is unbounded.

In 1987, Gyárfás [8] proposed to study families of graphs for which the chromatic number can be upper-bounded in terms of the clique number. More precisely, Gyárfás called a family of graphs \mathcal{G} *χ -bounded* if there is a function f such that $\chi(G) \leq f(\omega(G))$ for every $G \in \mathcal{G}$.

For a graph F , denote by $\text{Forb}(F)$ the family of graphs that contain no induced copy of F . This is a particularly interesting class of graphs for studying χ -boundedness, as it leads us to various examples and conjectures. Namely, one may ask: for which graphs F is $\text{Forb}(F)$ χ -bounded? If F contains a cycle, $\text{Forb}(F)$ is not χ -bounded; indeed, this follows from the existence of graphs with arbitrarily large chromatic number and girth (where the *girth* of a graph is the length of its shortest cycle), due to Erdős [6]. Gyárfás proved [8] that $\text{Forb}(F)$ is χ -bounded when F is a path

*Department of Mathematics, Hanyang University, Seoul, and Extremal Combinatorics and Probability Group, Institute for Basic Sciences (IBS), Daejeon, South Korea. E-mail: joonkyunglee@hanyang.ac.kr. Research supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government MSIT NRF-2022R1C1C1010300, Samsung STF Grant SSTF-BA2201-02, and IBS-R029-C4.

[†]Department of Mathematics, University College London, London WC1E 6BT, UK. Email: s.letzter@ucl.ac.uk. Research supported by the Royal Society.

[‡]Department of Mathematics, University College London, London WC1E 6BT, UK. Email: a.pokrovskiy@ucl.ac.uk.

or a star. An intriguing conjecture regarding χ -boundedness, due to Gyárfás [7] and Sumner [15], asserts that $\text{Forb}(F)$ is χ -bounded for every forest F . If true, this would solve the above question regarding graphs F for which $\text{Forb}(F)$ is χ -bounded. The conjecture is known for some special cases, including trees of radius 2 [9] and of radius 3 [10], but is widely open in general.

For a graph F , denote by $\text{Forb}^*(F)$ the family of graphs that do not contain an induced copy of a *subdivision* of F , where a subdivision of F is a graph obtained by replacing edges of F by internally disjoint paths. Scott [14] proved the following weakening of the Gyárfás–Sumner conjecture: $\text{Forb}^*(F)$ is χ -bounded for every forest F . He also conjectured that $\text{Forb}^*(F)$ is χ -bounded for every graph F , but this turned out to be false [4, 13]. At the best of our knowledge, there seems to be no conjectured answer to the question that asks for which graphs F is $\text{Forb}^*(F)$ χ -bounded.


The fact that $\text{Forb}^*(F) \subseteq \text{Forb}(F)$ for every graph F makes it natural to consider an ‘interpolation’ between the two classes to ask an analogous question. More precisely, fix an edge subset E of F and consider the class \mathcal{F} of graphs that contain no induced copy of a graph obtained from F by only subdividing the edges in E , while leaving the other edges unchanged. For example, taking F to be the graph obtained by adding a diagonal to a 4-cycle and letting E be set of edges in the cycle, the family \mathcal{F} obtained as above is the family of graphs that do not have a cycle with exactly one *chord*; given a graph G , a *chord* in a cycle C in G is an edge that joins two vertices in C which are not adjacent in C . Trotignon and Vušković [18] showed that this family of graphs with no cycle with exactly one chord is χ -bounded (in fact, they give a detailed description of the structure of such graphs). Inspired by this, Aboulker and Bousquet [1] suggested to study the family \mathcal{C}_k of graphs that do not contain a cycle with exactly k chords. They conjecture that \mathcal{C}_k is χ -bounded for every $k \geq 1$, and prove it for $k \in \{2, 3\}$. Note that [18] establishes the conjecture for $k = 1$. Our main result is to prove the Aboulker–Bousquet conjecture for sufficiently large k .

Theorem 1.1. *For every large enough k ,¹ there is a function f_k such that, if G is a graph with no cycle with exactly k chords, then $\chi(G) \leq f_k(\omega(G))$.*


A *k-wheel* is a cycle with an additional vertex that is adjacent to exactly k vertices in the cycle, and a *wheel* is a 3-wheel. It is easy to see that a k -wheel contains a cycle with exactly ℓ chords, for any $\ell \leq k - 2$ (see Figure 1).

It is thus quite natural, given our discussion of graphs with cycles with a given number of chords, to also consider graphs avoiding wheels. Trotignon [17] conjectured that the class of wheel-free graphs is χ -bounded. This conjecture is now known to be false (see Davies [5]), but a weaker version of it was proved by Bousquet and Thomassé [3]; they showed that, for every integer $\ell \geq 1$, the class of graphs with no induced wheel or an induced $K_{\ell, \ell}$ is χ -bounded. They also proved a similar result for k -wheels, with an additional triangle-free requirement; namely, they showed that, for every ℓ , the class of graphs with no induced k -wheel, $K_{\ell, \ell}$ or triangle is χ -bounded. Since a 2ℓ -cycle in $K_{\ell, \ell}$ has $\ell(\ell - 2)$ chords, taking $k = \ell(\ell - 2) + 2$ shows that the family of triangle-free graphs with no

¹Concretely, it suffices to take $k \geq 10^{14}$.



A 7-wheel



A cycle with four chords in a 7-wheel

Figure 1: Wheels

cycle with exactly $\ell(\ell - 2)$ chords is χ -bounded. We strengthen the latter statement by removing the triangle-free requirement.

Theorem 1.2. *For every integer $\ell \geq 3$ there is a function g_ℓ such that, if G is a graph with no cycles with exactly $\ell(\ell - 2)$ chords, then $\chi(G) \leq g_\ell(\omega(G))$.*

The proof of our main result, Theorem 1.1, is rather technical. We thus include also a proof of a weaker result, allowing us to find, given large k and a graph whose chromatic number is much larger than its clique number, a cycle with k' chords, where $k' \in \{k, k + 1, k + 2, k + 3\}$; see Theorem 6.1. This proof already includes a lot of the main ideas that come into the proof of the main, exact result, but avoid many of the technical difficulties.

In the next section, we give an outline of the proofs, focusing on the relatively simpler results Theorems 1.2 and 6.1. After that, in Section 3, we define the basic notions that we shall need and mention related observations. In Section 4 we show that, given an integer $k \geq 1$, graphs whose chromatic number is much larger than their clique number either contain a large induced bipartite graph or a cycle with exactly k chords. This will allow us to immediately deduce Theorem 1.2, and will also be a key component in the proofs of Theorems 1.1 and 6.1. In Section 5 we provide two number theoretic lemmas, related to Lagrange's four square theorem. Finally, in Section 6 we prove the approximate version of our main result, namely Theorem 6.1, and in Section 7 we prove the main result, Theorem 1.1.

2 Proof overview

The first step in our proof (accomplished in Section 4) proves the following: given positive integers ℓ and k , if G is a graph with large enough chromatic number, then it contains either an induced copy of $K_{\ell,\ell}$ or a cycle with exactly k chords (see Corollary 4.2). Our proof of this result partly follows the proof of a related result due to Bousquet and Thomassé [3], which asserts that, given ℓ and k , every graph with large enough chromatic number contains a triangle, an induced $K_{\ell,\ell}$, or a k -wheel. Recall that every $(k + 2)$ -wheel (defined before Figure 1) contains a cycle with exactly k chords (see Figure 1). Bousquet–Thomassé's result implies a weaker version of our first step, where

the graph G is assumed to be triangle-free. To avoid the triangle-free assumption, we leverage the fact that cycles with a prescribed number of chords are easier to find than wheels. This immediately implies our second main result, Theorem 1.2.

The next step connects to a classical result in number theory, which may be of independent interests. Recall that Lagrange’s four square theorem asserts that every natural number can be expressed as the sum of four integer squares. In Section 5, we prove two variants of this result. The first (Lemma 5.1) shows that every large enough integer can be expressed as the sum of exactly 20 integer squares that are all larger than a given constant. The second result, given in Lemma 5.2, is a similar statement, but here we aim to express a large number x as the sum of integers of the form $a(a + 1)$, for large a . To achieve this we require that x be divisible by 4 and we express it as a sum of 80 numbers of the aforementioned form.

In Section 6, we prove an approximate version (Theorem 6.1) of our main result where instead of finding a cycle with exactly k chords, we find a cycle whose number of chords is in $\{k, k + 1, k + 2, k + 3\}$. To do this, we first find, for some large ℓ , many induced copies of $K_{\ell, \ell}$ with no edges between them, by applying the first paragraph above several times. To find a cycle with the required number of chords, we take paths of appropriate lengths in the copies of $K_{\ell, \ell}$, and join these by *unimodal paths* (unimodal paths are a commonly used object in the study of χ -boundedness; we will explain what they are in the next section). A key point here is that the unimodal connections contribute $O(\sqrt{k})$ chords (see Lemmas 6.4 and 6.5). To get the right number of chords, we apply one of the two number-theoretic lemmas from the previous paragraph, depending on a certain parity condition. The case that requires the use of the second lemma above is the reason why this approach yields only an approximate result.

In Section 7 we prove the exact version of our main result, Theorem 1.1. To achieve this we proceed similarly to the paragraph above, but analyse much more carefully the interaction between the various copies of $K_{\ell, \ell}$; this leads to a rather technical proof. We will give an overview of this proof at the beginning of Section 7.

3 Preliminaries

Recall that a *chord* in a cycle C is an edge joining two non-consecutive vertices in C . Analogously, a *chord* in a path P is an edge that joins two non-consecutive vertices in P . It will be convenient to note that, given a graph G and a cycle C , the number of chords in C is $e(G[V(C)]) - |C|$.

We now introduce the notions of *extractions* and *unimodal paths*, which are commonly used in the study of χ -boundedness. The motivation behind these notions is the following observation. For a graph G , a vertex u and an integer $i \geq 0$, denote by $N_i(u)$ the set of vertices in G at distance exactly i from u . For brevity, given a set of vertices U , we write $\chi(U)$ to denote $\chi(G[U])$.

Observation 3.1. *Let u be a vertex in a graph G . Then there is an integer $i \geq 1$ such that $\chi(N_i(u)) \geq \chi(G)/2$.*

Proof. Let $U_0 = \bigcup_{i \geq 0} N_{2i}(u)$ and let $U_1 = \bigcup_{i \geq 0} N_{2i+1}(u)$. Since there are no edges between distinct $N_{2i}(u)$'s, $\chi(U_0) = \max_{i \geq 0} \chi(N_{2i}(u))$. In particular, there exists i_0 such that $\chi(U_0) = \chi(N_{2i_0}(u))$. By the same argument, there exists i_1 such that $\chi(U_1) = \chi(N_{2i_1+1}(u))$. Since $\{U_0, U_1\}$ is a partition of $V(G)$, we have $\chi(G) \leq \chi(U_0) + \chi(U_1) = \chi(N_{2i_0}(u)) + \chi(N_{2i_1+1}(u))$. Therefore, $\max\{\chi(N_{2i_0}(u)), \chi(N_{2i_1+1}(u))\} \geq \chi(G)/2$, as claimed. \square

Trivially, every vertex in $N_i(u)$ has a ‘backward’ neighbour in $N_{i-1}(u)$. Let $N_{\leq i}(u) := \bigcup_{0 \leq j \leq i} N_j(u)$. By chasing the backward neighbours of two vertices x and y in $N_i(u)$ repeatedly until they first intersect or have an edge between, we obtain a path between x and y in $N_{\leq i}(u)$ that is induced except the edge xy , if it is an edge in G . If this path has length ℓ , then the distance- j vertices, $j \leq \lfloor \ell/2 \rfloor$, from x or y are in $N_{i-j}(u)$.

Together with these paths, Observation 3.1 can be reformulated in a more systematic way. Namely, every graph G has a subgraph G_1 with the following properties:

- $\chi(G_1) \geq \chi(G)/2$,
- every vertex in G_1 has a neighbour in $V(G) \setminus V(G_1)$,
- for every distinct pair of vertices x and y in G_1 , there is a path P between x and y satisfying the following: the path is induced except for the edge xy ; the interior of P lies in $V(G) \setminus V(G_1)$; and the vertices in P apart from x, y , and their neighbours in P send no edges to $V(G_1)$.

Such a subgraph G_1 is called an *extraction* (or a *1-extraction*) of G , and the path joining two vertices in G_1 with the above properties is called a *unimodal path* with respect to G_1 .

One can repeat this process to obtain a sequence $G_0 = G \supseteq G_1 \supseteq \dots \supseteq G_p$ such that G_i is an extraction of G_{i-1} . Such a sequence $G_0 \supseteq \dots \supseteq G_p$ is called *extractions* and G_p is a *p-extraction* of G . The *i-th layer* of an extractions means $V(G_{i-1}) \setminus V(G_i)$. We will often say that G_p is a *p-extraction* of G without specifying the corresponding sequence of extractions. Once a *p-extraction* G_p of G is fixed, a *unimodal path in the i-th layer* is a unimodal path with respect to G_i with ends in G_p . This is an induced path except for the edges between the ends in G_p , whose interior is contained in $V(G_{i-1}) \setminus V(G_i)$. Moreover, the vertices in P apart from the end vertices and their neighbours in the path send no edges to $V(G_i)$. The following observation is immediate from the definitions, yet crucial for our arguments.

Observation 3.2. *Let P and Q be unimodal paths in a p -extraction of a graph G in layers i and j , respectively, with $i < j$. Then the vertices in P other than the end vertices and their neighbours in the path send no edges to $V(Q)$.*

As a final piece of notation, given an index $i \in [0, p]$ and a vertex x in G_i , an *i-father* of x is a neighbour of x in $V(G_{i-1}) \setminus V(G_i)$.

We close this section by mentioning the classical Kővari–Sós–Turán theorem, which we shall use frequently in what follows. The theorem will allow us to find large complete bipartite graphs in bipartite graphs with positive edge density.

Theorem 3.3 (Kővari–Sós–Turán). *For every $\varepsilon > 0$ and a positive integer ℓ , there exists n_0 such that the following holds: let G be a bipartite graph with a bipartition $A \cup B$ such that $|A|, |B| \geq n_0$. If G has at least $\varepsilon|A||B|$ edges, then G contains a copy of $K_{\ell, \ell}$.*

4 Obtaining large complete bipartite graphs

The main purpose of this section is to prove the following theorem, which finds ‘large’ complete bipartite graphs in graphs G with large chromatic number which have no cycle with exactly k chords.

Theorem 4.1. *For every integers $k, \ell \geq 1$, there exists a function g such that for every graph G one of the following holds: $\chi(G) \leq g(\omega(G))$; G contains a $K_{\ell, \ell}$; or G contains a cycle with exactly k chords.*

The following corollary of the previous theorem finds induced copies of $K_{\ell, \ell}$ in graphs G with large chromatic number and no cycle with k chords.

Corollary 4.2. *For every integers $k, \ell \geq 1$ there exists a function h such that for every graph G one of the following holds: $\chi(G) \leq h(\omega(G))$; G contains an induced $K_{\ell, \ell}$; or G contains a cycle with exactly k chords.*

Proof. Let $g_{k, \ell}$ be a function as in Theorem 4.1 for k and ℓ . For each $\omega \geq 1$, we define $h_{k, \ell}(\omega) := g_{k, L}(\omega)$ for some L satisfying $L \gg \ell, \omega$. Now consider a graph G . By choice of $h_{k, \ell}$, one of the following holds: $\chi(G) \leq g_{k, L}(\omega(G)) = h_{k, \ell}(\omega(G))$; G contains a $K_{L, L}$; or G contains a cycle with exactly k chords. In all but the second case, we are done, so suppose that G contains a $K_{L, L}$ and denote the corresponding bipartition by $\{A, B\}$. By Ramsey’s theorem and the choice of L , each of $G[A]$ and $G[B]$ contain an independent set of size ℓ . It follows that G contains an induced $K_{\ell, \ell}$, as required. \square

Notice that this corollary immediately implies Theorem 1.2, using the observation that an induced $K_{\ell, \ell}$ contains a cycle with exactly $\ell(\ell - 2)$ chords.

Before turning to the proof of Theorem 4.1, we mention two preliminary results. The first one, due to Thomas and Wollan, shows that $10k$ -connected graphs are k -linked, namely one can join any k pairs of vertices by pairwise vertex-disjoint paths. This improved on an earlier result of Bollobás and Thomason [2].

Theorem 4.3 ([16]). *Let $k \geq 0$ and let G be a $10k$ -connected graph. For every set of (not necessarily distinct) vertices $x_1, \dots, x_k, y_1, \dots, y_k$ there exist paths Q_1, \dots, Q_k with pairwise disjoint interiors, such that Q_i has ends x_i, y_i .*

The second preliminary result, due to Kühn and Osthus [11], allows us to find a subdivision of a κ -connected graph.

Theorem 4.4. *Let $\chi \gg \kappa, \ell$. Then for every graph G one of the following holds: $\chi(G) \leq \chi$; G contains a $K_{\ell, \ell}$; or G contains, as an induced subgraph, a 1-subdivision of a κ -connected graph H .*

Recall that a 1-subdivision of a graph H is the graph obtained from H by replacing each edge uv by a path $uw_{uv}v$, where the vertices w_{uv} are distinct. For brevity, denote by H_{sb} the 1-subdivision of a graph H .

4.1 Finding a large complete bipartite graph

In the remainder of this section, we prove Theorem 4.1. To do so, we assume that we are given a graph G with large chromatic number and which contains no large complete bipartite subgraphs. The starting is Theorem 4.4 that says that under this assumption G contains an induced copy of a 1-subdivision of a highly connected graph H . In fact, we may assume that this subdivision lies in a p -extraction of G , for some large p . Our proof splits-off into two cases: either we can emulate the triangle-freeness assumption that Bousquet and Thomassé [3] impose (when proving that every triangle-free graph with large chromatic number contains either a large complete bipartite subgraph or a long wheel), or not. In the former case our proof largely follows [3], and in the latter case we use the abundance of triangles to construct a cycle with the right number of chords.

Throughout this section, we use the following setup. The parameters are chosen according to the hierarchy

$$\chi \gg p \gg \kappa \gg k_1 \gg k_2 \gg k, \ell, \omega.$$

Let G be a $K_{\ell, \ell}$ -free graph with $\chi(G) = \chi$ and $\omega(G) \leq \omega$. Let $G \supseteq G_1 \supseteq \dots \supseteq G_p$ be a sequence of extractions, so that $\chi(G_p) \geq 2^{-p}\chi$. We will show that G contains a cycle with exactly k chords.

By Theorem 4.4, there is a κ -connected graph H whose 1-subdivision H_{sb} is an induced subgraph of G_p . Fix an induced copy of H_{sb} in G_p and denote by $\text{sub}(e)$ the vertex of the induced copy of H_{sb} in G_p that subdivides the edge e in H . We also identify vertices in H as corresponding vertices in the copy of H_{sb} (and hence, in G_p). When considering a neighbour z of a vertex x in H , we say that z is an H -neighbour of x to stress that the adjacency is not in the host graph G_p .

The following lemma allows us to deal with the case where at every vertex x in H , for many H -neighbours z of x , $\text{sub}(xz)$ has a common j -father with at least one of x and z .

Lemma 4.5. *Suppose that every vertex x in H is incident with at least $2k_1$ edges $e = xy$ in H such that $\text{sub}(e)$ has a common j -father with at least one of x and y , for at least k_1 indices j . Then G contains a cycle with exactly k chords.*

We postpone the proof of this lemma until the end of this section. In the light of Lemma 4.5, we may assume that the particular structure described therein does not appear in the fixed copy of H_{sb} in G_p . That is, there is a ‘special’ vertex x in H such that for all but at most $2k_1$ neighbours z of x in H the following holds: for all but at most k_1 values of j , the vertex $\text{sub}(xz)$ does not have a common j -father with neither x nor z . Let Z be a set of k_1 neighbours of x in H that satisfy this property, i.e., for $z \in Z$, there are at most $2k_1$ indices j such that $\text{sub}(xz)$ has a common j -father with x or z .

Then all but at most $2k_1^2$ values of j satisfy: for every $z \in Z$, the vertex $\text{sub}(xz)$ does not have a common j -father with neither x nor z . Let J be the set of ‘good’ indices j that satisfy this property. In particular, $|J| \geq p - 2k_1^2$. The vertex x , set $Z \subseteq N_H(x)$ and set $J \subseteq [p]$ will be fixed throughout this section. Let Y be the set of vertices in the fixed copy of H_{sb} in G_p that are adjacent to x and some vertex in Z , i.e., $Y := \{\text{sub}(xz) : z \in Z\}$. For $y \in Y$, the *extended neighbour* $\text{ext}(y)$ of y is the vertex $z \in Z$ such that $y = \text{sub}(xz)$. In particular, $|Z| = |Y| = k_1$.

Following the definition in page 7 of [3], a collection of unimodal paths \mathcal{Q} with the set $Y' \subseteq Y$ of endpoints is α -good for $\alpha > 0$ if the following conditions hold:

- (G1) For every $y \in Y'$, there exists a unique path $Q \in \mathcal{Q}$ with the endpoint y and moreover, Q is the only path that contains an edge incident with y .
- (G2) For every $y \in Y'$, no vertex in any of the paths in \mathcal{Q} contains a neighbour of the extended neighbour $\text{ext}(y)$ of y other than y itself.
- (G3) For any distinct $Q, Q' \in \mathcal{Q}$, there is no edge between a father (in Q) of an endpoint in Q and a father (in Q') of an endpoint in Q' .
- (G4) For every $y \in Y'$ and every $Q \in \mathcal{Q}$, there are at most $\alpha |N_{H_{\text{sb}}}(\text{ext}(y))|$ vertices in $N_{H_{\text{sb}}}(\text{ext}(y))$ that are adjacent to a vertex in Q .

We say that a collection of unimodal paths \mathcal{Q} is *independent* if there are no edges between distinct paths in \mathcal{Q} . The following is a variant of the first part of (the proof of) Lemma 9 in [3].

Lemma 4.6. *There is a $\frac{1}{4k_2}$ -good collection of unimodal paths \mathcal{Q} of size at least k_2 in G .*

Proof. Let k', k'', c be such that $k_1 \gg k' \gg k'' \gg c \gg k_2$. For each $y \in Y$ and $j \in J$, choose a j -father of y and denote it by $f_j(y)$. We wish to find ‘large’ subsets $Y' \subseteq Y$ and $J' \subseteq J$ such that, for every fixed j , the fathers $f_j(y)$ are distinct for $y \in Y'$. To this end, we construct sequences $Y = Y_0 \supseteq \dots \supseteq Y_\ell$ and $j_1, \dots, j_\ell \in J$ recursively as follows: given Y_0, \dots, Y_t and j_1, \dots, j_t for $t < \ell$, if there exists $j \in J \setminus \{j_1, \dots, j_t\}$ and a subset $Y' \subseteq Y_t$ of size at least $\sqrt{|Y_t|}$ such that $f_j(y)$ is the same for all $y \in Y'$, define $j_{t+1} = j$ and $Y_{t+1} = Y'$. Otherwise, stop the process.

Suppose first that j_1, \dots, j_ℓ and Y_ℓ are well-defined through the recursive process. Then $f_j(y)$ is the same for all $y \in Y_\ell$, for every $j \in \{j_1, \dots, j_\ell\}$. Then $\{f_j(y) : j \in \{j_1, \dots, j_\ell\}\}$ is a set of size ℓ that

is fully joined to Y_ℓ . Since $|Y_\ell| \geq |Y|^{2^{-\ell}} = (k_1)^{2^{-\ell}} \geq \ell$, there is a copy of $K_{\ell,\ell}$ in G , contradicting the assumption that G is $K_{\ell,\ell}$ -free.

Now suppose that the process stops at the t -th step for some $t < \ell$ with the outputs j_1, \dots, j_t and Y_t . Let $J' := J \setminus \{j_1, \dots, j_t\}$. Then for every $j \in J'$, each element in the multiset $\{f_j(y) : y \in Y_t\}$ repeats at most $\sqrt{|Y_t|}$ times. Since $|Y_t| \geq |Y|^{2^{-t}} = (k_1)^{2^{-t}} \geq (k'')^2$, for every $j \in J'$ there is a subset $Y'_j \subseteq Y_t$ of size k'' such that $f_j(y)$ are distinct for all $y \in Y'_j$. Let $Y' \subseteq Y_t$ be the most popular choice for Y'_j . By averaging, $Y'_j = Y'$ for at least $|J'| / \binom{|Y_t|}{\sqrt{|Y_t|}} \geq (p - k_1^2 - \ell) / 2^{k_1} \geq k'$ indices $j \in J'$; let $J'' \subseteq J'$ be a set of size k' such that $Y'_j = Y'$ for every $j \in J''$.

Let $W := \{f_j(y) : y \in Y', j \in J''\}$. We claim that for every $y \in Y'$ there are at most k'' elements $w \in W$ such that w sends at least $\frac{1}{4k_2} |N_{H_{\text{sb}}}(\text{ext}(y))|$ edges to $N_{H_{\text{sb}}}(\text{ext}(y))$. Suppose to the contrary that this is not the case for $y \in Y'$. Write $N := N_{H_{\text{sb}}}(\text{ext}(y))$ for brevity and let W' be the set of vertices $w \in W$ such that w sends at least $\frac{1}{4k_2} |N|$ edges to N satisfying $|W'| \geq k''$. By Theorem 3.3, $G[N, W']^2$ contains a copy of $K_{\ell,\ell}$, a contradiction. Consider the set

$$\left\{ j \in J'' : \text{for all } z, y \in Y', f_j(z) \text{ sends at most } \frac{1}{4k_2} |N_{H_{\text{sb}}}(\text{ext}(y))| \text{ edges to } N_{H_{\text{sb}}}(\text{ext}(y)) \right\}.$$

By the above statement, this set has size at least $|J''| - |Y'| \cdot k'' \geq k' - (k'')^2 \geq k_2$. Let J_{final} be a subset of the above set of size k_2 .

For every $j \in J_{\text{final}}$, define a graph F_j on vertices Y' whose edges are pairs $y_1 y_2$ where $y_1, y_2 \in Y'$ are distinct and one of the following holds: $f_j(y_i)$ is adjacent to y_{3-i} for some $i \in [2]$; $f_j(y_i)$ is adjacent to $\text{ext}(y_{3-i})$; or $f_j(y_1)$ and $f_j(y_2)$ are adjacent. By Ramsey's theorem, the graph $F = \bigcup_{j \in J_{\text{final}}} F_j$ contains either an independent set of size $2k_2$ or a clique of size c . Assume the latter case and suppose that U is a subset of Y' of size c that induces a clique. Then for some $j \in J_{\text{final}}$ the graph F_j has at least $\frac{1}{k_2} \binom{|U|}{2}$ edges. Let $U' := U \cup \{\text{ext}(y) : y \in U\} \cup \{f_j(y) : y \in U\}$. Then, by definition of F_j ,

$$e(G[U']) \geq e(F_j) \geq \frac{1}{k_2} \binom{|U|}{2} \geq \frac{1}{k_2} \binom{|U'|/3}{2}.$$

By Theorem 3.3, $G[U']$ contains a copy of $K_{\ell,\ell}$, a contradiction.

It remains to consider the case where F contains an independent set U of size $2k_2$. Write $U = \{y_1, \dots, y_{2k_2}\}$ and $J_{\text{final}} = \{j_1, \dots, j_{k_2}\}$. Let Q_i be a unimodal path between $f_{j_i}(y_{2i-1})$ and $f_{j_i}(y_{2i})$, for $i \in [k_2]$. By construction, one may easily check that $\{Q_1, \dots, Q_{k_2}\}$ is a $\frac{1}{4k_2}$ -good collection of size at least k_2 . \square

Recall that a k -wheel in a graph F is an induced cycle C along with an additional vertex that has at least k neighbours in C . We note that a $(k+2)$ -wheel contains a cycle with exactly k chords. Indeed, suppose that C is an induced cycle and v is a vertex (not in C) with at least $k+2$ neighbours in C . Let u_0, \dots, u_{k+1} be $k+2$ consecutive neighbours of v in C . Let P be the subpath in C that

²Here $G[A, B]$ denotes the *bipartite* induced subgraph, i.e., we only take those edges that cross between A and B .

starts at u_0 , ends in u_{k+1} and contains the vertices u_1, \dots, u_k , and let C' be the cycle obtained by concatenating P and the path $u_{k+1}vu_0$. Then C' is a cycle with exactly k chords; its chords are vu_1, \dots, vu_k .

The following lemma obtains a collection of independent good paths from a collection of good paths. We use this as a black box.

Lemma 4.7 (Lemma 17 in [3]). *Let \mathcal{Q} be an α -good collection of unimodal paths of size k_2 . If G is k -wheel-free, then there exists an independent 2α -good collection of unimodal paths of size at least k .*

The following lemma, together with the lemmas above, will complete the proof of Theorem 4.1.

Lemma 4.8 (Lemma 12 in [3]). *If there is a collection of $\lceil k/2 \rceil$ independent $(1/2k)$ -good unimodal paths with endpoints in $N_{H_{\text{sb}}}(x)$ for a vertex x in H , then the graph G has a k -wheel.*

Proof of Theorem 4.1. Recall that the parameters are chosen according to the hierarchy $\chi \gg p \gg \kappa \gg k_1 \gg k_2 \gg k, \ell, \omega$. By Lemma 4.6, there is a collection \mathcal{Q} that consists of k_2 paths that are $\frac{1}{4k_2}$ -good. Then by Lemma 4.7, there is a collection of k_2 independent $\frac{1}{2k_2}$ -good paths; however, Lemma 4.8 then finds a k_2 -wheel in G , and hence, a cycle with k chords. \square

4.2 Dealing with triangles

It remains to prove Lemma 4.5. The following lemma will be useful in the proof.

Lemma 4.9. *Suppose that G does not have a cycle with exactly k chords. Then for every vertex u in G , there exists a subset $R \subseteq V(H)$ of at most $3(k+1)$ vertices such that u does not have any edges to the copy of $(H \setminus R)_{\text{sb}}$ in G .*

Proof. Let X be the set of vertices in H that are neighbours of u in G . We claim that $|X| \leq k+1$. Suppose to the contrary that x_1, \dots, x_{k+2} are distinct vertices in X . As H is κ -connected with $\kappa \gg k$, Theorem 4.3 guarantees that there exist pairwise internally vertex-disjoint paths Q_1, \dots, Q_{k+2} such that each Q_i ends at x_i and x_{i+1} , where the addition of indices taken modulo $k+2$. Let Q'_i be the path in the copy of H_{sb} in G that corresponds to the 1-subdivision of Q_i . Then by concatenating the paths Q'_1, \dots, Q'_{k+2} , we obtain an induced cycle in G that contains at least $k+2$ neighbours of u . Then there exists a cycle with exactly k chords, which contradicts the assumption.

Let M be a maximum matching in H such that, for each edge e in M , $\text{sub}(e)$ is adjacent to u in G . Our next claim is $|M| \leq k+1$. Suppose to the contrary that $x_1y_1, \dots, x_{k+2}y_{k+2}$ are vertex-disjoint edges in M . By using Theorem 4.3 as before, there exist vertex-disjoint paths Q_1, \dots, Q_k in $H \setminus \{x_1y_1, \dots, x_{k+2}y_{k+2}\}$ such that each Q_i ends at y_i and x_{i+1} . Again, let Q'_i be the 1-subdivision of Q_i in the copy of H_{sb} in G . Then $x_1y_1Q'_1 \dots x_{k+2}y_{k+2}Q'_{k+2}x_1$ is an induced cycle in G that contains at least $k+2$ neighbours of u , which again yields a contradiction.

Take $R = X \cup V(M)$. Then $|R| \leq 3(k+1)$ and u has no neighbours in $(H \setminus R)_{\text{sb}}$, as required. \square

Proof of Lemma 4.5. Let k_3, k_4 be such that $k_1 \gg k_2 \gg k_3 \gg k_4 \gg \omega \gg k$. Say that an edge $e = xy$ of H is *triangulated* if $\text{sub}(e)$ and at least one of x and y have a common j -father, for at least k_1 indices j . Then every vertex in H is incident with at least $2k_1$ triangulated edges and hence, there is a matching M_1 in H that consists of k_1 triangulated edges.

For each $e \in M_1$ let $x(e)$ and $y(e)$ be the two ends of e , let $f(e)$ be a $j(e)$ -father of $\text{sub}(e)$ which is adjacent to $x(e)$ or $y(e)$, so that the indices $j(e)$ are all distinct for $e \in M_1$. Let U be the set $\{f(e) : e \in M_1\}$; then $|U| = k_1$. Since $k_1 \gg k_2, \omega$ and G is $K_{\omega+1}$ -free, the set U contains an independent set of size k_2 . Let M_2 be a submatching of M_1 of size k_2 such that $\{f(e) : e \in M_2\}$ is independent.

Let F be the auxiliary graph whose vertices are the edges of M_2 , where $e_1 e_2$ is an edge in F whenever $f(e_{3-i})$ is adjacent to at least one of $\text{sub}(e_i), x(e_i)$ and $y(e_i)$ for some $i \in [2]$. By Lemma 4.9, the vertex $\text{sub}(e)$ is adjacent to at most $3(k+1)$ vertices in the copy of H_{sb} in G , for every $e \in M_2$. In particular, every $e \in M_2$ has neighbours in at most $3(k+1)$ of the sets $\{\text{sub}(e'), x(e'), y(e')\}$ with $e' \in M_2$. Thus, F has at most $3(k+1)|F|$ edges. Turán's theorem then implies that F contains an independent set of size at least $|F|/6(k+1) \geq k_3$. Let M_3 be a submatching of M_2 of size k_3 that forms an independent set in F . That is, for every distinct $e, e' \in M_3$, $f(e)$ is *not* adjacent to any $\text{sub}(e'), x(e')$ or $y(e')$.

For each $e \in M_3$, let $R(e)$ be a set of vertices in H of size at most $3(k+1)$ such that $x(e'), y(e') \notin R(e)$ for all $e' \in M_3$ and $f(e)$ has no neighbours in $(H \setminus R(e))_{\text{sb}}$ except for possibly $\text{sub}(e), x(e)$ and $y(e)$; the existence of such a set is guaranteed by the choice of M_3 and by Lemma 4.9.

We consider three cases. Throughout the case analysis, we abuse notation by writing e for the vertex $\text{sub}(e)$ for simplicity. Whenever we consider indexed edges e_1, e_2, \dots , we shall denote $x_i := x(e_i), y_i := y(e_i), f_i := f(e_i)$ and $R_i := R(e_i)$. The 1-subdivision of a path Q_i in H is denoted by Q'_i , and is again a path in the copy of H_{sb} .

Case 1: there are k edges $e \in M_3$ such that $f(e)$ is not adjacent to $y(e)$.

Let $e_1, \dots, e_k \in M_3$ be distinct edges such that f_i is not adjacent to y_i for $i \in [k]$ (by choice of M_3 this means that f_i is adjacent to x_i for $i \in [k]$). Let $R := R_1 \cup \dots \cup R_k$. As H is κ -connected, $H \setminus R$ is $10k$ -connected and thus, by Theorem 4.3, there exist pairwise vertex-disjoint paths Q_1, \dots, Q_k in $H \setminus R$ such that Q_i has ends y_i and x_{i+1} for $i \in [k]$, where the addition of indices is taken modulo k . Let C be the cycle $x_1 f_1 e_1 y_1 Q'_1 \dots x_k f_k e_k Q'_k x_1$. Then C has exactly k chords, namely: $x_1 e_1, \dots, x_k e_k$.

Figure 2: Case 1: f_i not adjacent to y_i

Case 2: k is even and $f(e)$ is adjacent to $x(e)$ and $y(e)$ for at least $k/2$ values of e in M_3 .

Write $k = 2s$. Let $e_1, \dots, e_s \in M_3$ be distinct edges such that f_i is adjacent to x_i and y_i for $i \in [s]$. Let $R := R_1 \cup \dots \cup R_s$. As before, there exist pairwise vertex-disjoint paths Q_1, \dots, Q_s in $H \setminus R$

such that Q_i has ends y_i and x_{i+1} for $i \in [s]$, where the addition of indices is taken modulo s . Then $x_1 f_1 e_1 y_1 Q'_1 \dots x_s f_s e_s y_s Q'_s x_1$ is a cycle in G with exactly k chords: $x_1 e_1, \dots, x_s e_s$ and $f_1 y_1, \dots, f_s e_s$.

Figure 3: Case 2: k even f_i adjacent to x_i and y_i

Case 3: k is odd and $f(e)$ is adjacent to $x(e)$ and $y(e)$ for at least $k_4 + 1$ values of e in M_3 .

Note that if $k = 1$ then $f(e)x(e)ey(e)$ is a cycle with exactly one chord, for any $e \in M_3$ where $f(e)$ is adjacent to $x(e)$ and $y(e)$. We may thus assume that $k \geq 3$; write $k = 2s - 1$ with $s > 1$.

Let M_4 be a submatching of M_3 of size k_4 that consists of edges e where $f(e)$ is adjacent to $y(e)$ and $j(e) > 1$ (recall that M_3 contains at most one edge e with $j(e) = 1$). For each $e \in M_4$, let $g(e)$ be a 1-father of $f(e)$. Here the $g(e)$'s are not necessarily distinct. We claim that each $g(e)$, with $e \in M_4$, is adjacent to at most $k + 1$ of the vertices $f(e')$ with $e' \in M_4$. Indeed, suppose that e_1, \dots, e_{k+2} are distinct edges in M_4 such that each f_j , with $j \in [s]$, is adjacent to $g(e)$. As usual, we can find paths Q_1, \dots, Q_{k+2} in $H \setminus (R(e_1) \cup \dots \cup R(e_{k+2}))$ that are pairwise vertex-disjoint and Q_i has ends y_i and x_{i+1} for $i \in [k+2]$. Then $x_1 f_1 y_1 Q'_1 \dots x_{k+2} f_{k+2} y_{k+2} Q'_{k+2} x_1$ is an induced cycle in G that contains at least $k + 2$ neighbours of $g(e)$. Thus, a cycle with exactly k chords exists.

Let F be the auxiliary graph on the edges in M_4 , where e_1 and e_2 form an edge if $g(e_{3-i})$ is adjacent to at least one of $e_i, x(e_i), y(e_i)$, and $f(e_i)$ for some $i \in [2]$. By Lemma 4.9 and the previous paragraph, each $g(e)$, with $e \in M_4$, is adjacent to at most $4k + 4$ vertices amongst $\{e', x(e'), y(e'), f(e') : e' \in M_4\}$. Therefore, F has at most $(4k + 4)|F|$ edges and thus, it has an independent set of size at least $|F|/(8k + 4) \geq k_4/(8k + 4) \geq \lceil k/2 \rceil$. Let e_1, \dots, e_s be distinct edges in M_1 that form an independent set in F . Write $g_i := g(e_i)$ and let R'_i be a set of at most $3(k + 1)$ vertices in H such that g_i has no neighbours in $(H \setminus R'_i)_{\text{sb}}$ except for possibly x_i, y_i, e_i , and $x_j, y_j, e_j \notin R'_i$ for $j \in [s]$. Such a set R'_i exists by Lemma 4.9 and the fact that e_i 's form an independent set in F . Observe that the g_i 's are distinct (since g_i is adjacent to f_i but not to f_j with $j \in [s] \setminus \{i\}$), and g_i sends no edges to $\{x_j, y_j, e_j, f_j\}$ for $j \neq i$. Let $R = R_1 \cup \dots \cup R_s \cup R'_1 \cup \dots \cup R'_s$. As usual, let Q_1, \dots, Q_{s-1} be pairwise vertex-disjoint paths in $H \setminus R$ such that each Q_i ends at y_i and x_{i+1} . Then the path $y_1 Q'_1 x_2 f_2 e_2 y_2 Q'_2 \dots x_{s-1} f_{s-1} e_{s-1} y_{s-1} Q_{s-1} x_s$ has exactly $2(s - 2) = k - 3$ chords. Similarly, paths with exactly $k - 3$ chords exist between any $u \in \{x_1, y_1\}$ and $v \in \{x_s, y_s\}$. To extend one such path to a cycle with exactly k chords, we need the following claim.

Claim 4.10. *There exist paths P_1 and P_s such that P_i has ends g_i and one of x_i, y_i and $V(P_i) \subseteq \{x_i, y_i, e_i, f_i, g_i\}$, for $i \in \{1, s\}$, and P_1 and P_s have three chords in total.*

Proof of the claim. Write $x = x_i, y = y_i, e = e_i, f = f_i, g = g_i$ for some $i \in \{1, s\}$, and let σ be the number of neighbours of g in $\{x, y, e\}$. We aim to show that there are paths P with vertices in $\{x, y, e, f, g\}$ whose ends are g and one of x, y satisfying the following requirements (separately, namely P varies).

- (0) P has no chords, if $\sigma = 1$;

- (1) P has exactly one chord, if $\sigma \neq 1$;
- (2) P has exactly two chords;
- (3) P has three chords, if $\sigma = 1$.

By using this, Claim 4.10 easily follows. Indeed, unless $\sigma = 1$ for both $i = 1$ and s , we can take one of P_1 and P_s to have one and two chords, respectively, using (2) and (1). Otherwise, we can take one to have no chords and the other to have three, using (0) and (3). To show that paths P that satisfies (1)–(4) exist, we consider the four possible values of σ .




Figure 4: Proof of Claim 4.10

- $\sigma = 0$. For (1) take $P = gfey$, and for (2) take $P = gfxy$.
- $\sigma = 1$. Without loss of generality, g is not adjacent to x . For (0), (2), and (3), take $P = gfx$, $P = gfey$, and $P = gfxy$, respectively.
- $\sigma = 2$. Without loss of generality, g is adjacent to x . For (1) and (2), take $P = gxy$ and $P = gfey$, respectively.
- $\sigma = 3$. For (1) and (2), take $P = gey$ and $P = gxy$, respectively. □

Let P_1 and P_s be as in Claim 4.10. Without loss of generality, we may assume that y_1 is an end of P_1 and x_s is an end of P_s . Let P be a unimodal path with ends g_1 and g_s . Then the interior of P sends no edges to the copy of H_{sb} or $\{f_1, \dots, f_s\}$, since each g_i is a 1-father of f_i 's, whereas the f_i 's and the copy of H_{sb} are in G_2 . Finally, augmenting the path $y_1Q'_1x_2f_2e_2y_2Q'_2 \dots x_{s-1}f_{s-1}e_{s-1}y_{s-1}Q'_{s-1}x_s$ by adding the path $x_sP_sg_sPg_1P_1y_1$ gives a cycle with exactly k chords. □

lem5case3.pdf

Figure 5: Case 3: odd k and f_i adjacent to x_i and y_i

5 Number theory lemmas

In this section we prove two results which are variants Lagrange's four-square theorem, which asserts that every positive integer can be written as the sum of at most four integer squares.

Lemma 5.1. *For every c , every large enough k can be written as a sum of exactly 20 squares larger than c^2 .*

Proof. Given a non-negative integer x , let $f(x) = x^2 + 2500c^2 + (4c + 1)^2$. We first claim that for every integer $x \geq 0$, the number $f(x)$ can be written as the sum of exactly five squares larger than c^2 . If $x \geq c$, this follows by writing

$$f(x) = x^2 + (30c)^2 + (24c)^2 + (32c)^2 + (4c + 1)^2.$$

If $x \leq c$ and x is even, we have

$$f(x) = 2 \left(\frac{50c + x}{2} \right)^2 + 2 \left(\frac{50c - x}{2} \right)^2 + (4c + 1)^2,$$

which readily implies that $f(x)$ is the sum of five squares larger than c^2 . Finally, if $x \leq c$ and x is odd, we use the following equality to reach the same conclusion.

$$x^2 + (4c + 1)^2 = 2 \left(\frac{4c + 1 + x}{2} \right)^2 + 2 \left(\frac{4c + 1 - x}{2} \right)^2.$$

Let $\ell := k - 4(2500c^2 + (4c + 1)^2)$ and suppose that k is large enough so that $\ell \geq 0$. By the four squares theorem, there exist non-negative integers x_1, \dots, x_4 such that $\ell = x_1^2 + x_2^2 + x_3^2 + x_4^2$. Equivalently,

$$k = \sum_{i=1}^4 (x_i^2 + 2500c^2 + (4c + 1)^2) = \sum_{i=1}^4 f(x_i).$$

By using the fact that each $f(x_i)$ is the sum of five squares larger than c^2 , we conclude that k is the sum of 20 squares larger than c^2 , as required. \square

Lemma 5.2. *For every c , for every large enough k which is divisible by 4, there exist $a_1, \dots, a_{80} \geq c$ such that $k = \sum_{i \in [80]} a_i(a_i + 1)$.*

Proof. By Lemma 5.1, the integer $k/4$ can be written as the sum of twenty squares larger than $(c+1)^2$. Write $k/4 = \sum_{i=1}^{20} x_i^2$, where $x_i \geq c+1$. Then $k = \sum_{i=1}^{20} 4x_i^2$, i.e., k is the sum of 20 even squares larger than $4(c+1)^2$. Observe now that every even square larger than $4(c+1)^2$ can be written as a sum $\sum_{i=1}^4 a_i(a_i+1)$ with $a_i \geq c$, as $4a^2 = 2 \cdot (a(a+1) + a(a-1))$. Thus, k is a sum $\sum_{i=1}^{80} a_i(a_i+1)$ with $a_i \geq c$, as required. \square

6 Proof of approximate result

The aim in this section is to prove the following result, that allows us to find a cycle of almost the required number of chords, in any graph whose chromatic number is much larger than the clique number.

Theorem 6.1. *Let k be large enough. Then there is a function f such that for every graph G either $\chi(G) \leq f(\omega(G))$ or G has a cycle with exactly k' chords for some $k' \in \{k, k+1, k+2, k+3\}$.*

The proof relies on results from previous sections, as well as the following two lemmas. The next lemma allows us to assume that there is a large collection of pairwise disjoint large induced balanced bipartite subgraphs with no edges between them.

Lemma 6.2. *Every graph G contains either ℓ pairwise vertex-disjoint induced copies of $K_{\ell, \ell}$ with no edges between them or an induced subgraph H with $\chi(H) \geq \chi(G)/(2\ell^2) - 1$ such that either $\omega(H) < \omega(G)$ or it is induced $K_{\ell, \ell}$ -free.*

Proof. Let K_1, \dots, K_t be a maximal collection of pairwise vertex-disjoint induced copies of $K_{\ell, \ell}$ with no edges between them; write $K = V(K_1 \cup \dots \cup K_t)$. We are done if $t \geq \ell$, so suppose that $t < \ell$. Notice that one of the subgraphs $G[N(v) \cup \{v\}]$ for some $v \in K$ or the graph $G \setminus (K \cup N(K))$ has chromatic number at least $\chi(G)/(2\ell^2)$, since the union of these $2\ell t + 1$ graphs covers all the vertices in G and $2\ell t + 1 \leq 2\ell^2$.

Suppose first that some $G[N(v) \cup \{v\}]$ has chromatic number at least $\chi(G)/(2\ell^2)$. Then $\chi(G[N(v)])$ is at least $\chi(G)/(2\ell^2) - 1$ and $\omega(G[N(v)]) < \omega(G)$, so we can take $H := G[N(v)]$. Otherwise, if $G \setminus (K \cup N(K))$ has chromatic number at least $\chi(G)/(2\ell^2)$, then $H := G \setminus (K \cup N(K))$ is induced $K_{\ell, \ell}$ -free subgraph by maximality of t . \square

The next lemma is the key ingredient in proving Theorem 6.1, whose proof will be given in Section 6.2.

Lemma 6.3. *Let $\ell \gg k \gg 1$,³ and let $p > 300$. In a p -extraction of a graph G , suppose that there are 101 induced copies of $K_{\ell, \ell}$ with no edges between them. Then there is a cycle with k' chords, for some $k' \in \{k, k+1, k+2, k+3\}$.*

³concretely, it suffices to take $k \geq 10^{12}$.

We now prove Theorem 6.1 by using the previous lemmas.

Proof of Theorem 6.1. We prove by induction on ω that there exists $f(\omega)$ such that, for a graph G with clique number ω , if $\chi(G) > f(\omega)$ then G has a cycle with exactly k' chords for some $k' \in \{k, k+1, k+2, k+3\}$. For $\omega = 1$, we can take $f(1) = 2$, for which the statement is vacuously true as there are no graphs with clique number 1 and chromatic number at least 2.

Suppose that for $\omega_0 \geq 1$, $f(\omega)$ is defined for $\omega \leq \omega_0$ so that the above statement holds. Let p and ℓ be sufficiently large, e.g., $p \geq 300$ and $\ell \geq 2^{500}\sqrt{k}$, and let g be a function as in Corollary 4.2 for k and ℓ ; namely, if $\chi(G) > g(\omega(G))$ then G contains either an induced $K_{\ell,\ell}$ or a cycle with exactly k chords. Now set $f(\omega_0 + 1) = 2^{p+1}\ell^2 \cdot (\max\{g(\omega_0), f(\omega_0)\} + 1)$. Consider a graph G with $\omega(G) = \omega_0 + 1$ and $\chi(G) > f(\omega_0 + 1)$. Our goal is to show that G contains a cycle with exactly k' chords, for some $k' \in \{k, k+1, k+2, k+3\}$. Let G' be a p -extracted graph of G . In particular, $\chi(G') \geq \chi(G)/2^p$. By Lemma 6.2, one of the following three cases holds for G' .

- (G1) there are ℓ pairwise vertex-disjoint induced copies of $K_{\ell,\ell}$ with no edges between them,
- (G2) there is an induced subgraph $H \subseteq G'$ with $\chi(H) \geq \chi(G')/(2\ell^2) - 1$ and $\omega(H) < \omega(G')$,
- (G3) there is an induced subgraph $H \subseteq G'$ with $\chi(H) \geq \chi(G')/(2\ell^2) - 1$ which is $K_{\ell,\ell}$ -free.

If (G2) holds, then $\chi(H) \geq \chi(G)/(2^{p+1}\ell^2) - 1 > f(\omega_0)$ and $\omega(H) \leq \omega_0$. Thus, by the inductive hypothesis, H contains a cycle with exactly k' chords, for some $k' \in \{k, k+1, k+2, k+3\}$.

If (G3) holds, then $\chi(H) \geq \chi(G)/(2^{p+1}\ell^2) - 1 > g(\omega_0)$. Since H is $K_{\ell,\ell}$ -free in this case, by choice of g there is a cycle with exactly k chords in H .

We may now assume that (G1) holds. In particular, there are 101 copies of $K_{\ell,\ell}$ in H that are pairwise vertex-disjoint with no edges between them. By Lemma 6.3, using the choice of G' as a p -extraction of G , there is a cycle in G with exactly k' chords, with $k \leq k' \leq k+3$. \square

6.1 Few edges between unimodal paths

For the proof of Lemma 6.3, we need the following two lemmas that allow us to assume that there are only few edges between two unimodal paths connected to a large induced balanced bipartite subgraph.

Lemma 6.4. *Let $\ell \gg k$, $p \geq 1$, and let K be an induced copy of $K_{\ell,\ell}$ in a p -extraction of a graph G . If P is a unimodal path starting in K and u is a vertex outside of P with an edge into $K \setminus V(P)$, then either there is a cycle in G with exactly k chords or there are at most $8\sqrt{k}$ edges from u to P .*

Proof. Let x and y be the ends of P with $x \in V(K)$ and let x^- and y^- be the unique neighbours of x and y in P . By unimodality, vertices in K do not send any edges to $P \setminus \{x, x^-, y, y^-\}$. Let w be a neighbour of u in K distinct from x , which exists by the assumption on u .




Figure 6: Lemma 6.4

Suppose that u sends more than $8\sqrt{k}$ edges to P . Let Q be a path in K with ends x and w on $2a$ or $2a + 1$ vertices, where $a = \lfloor \sqrt{k} - 2 \rfloor$. Let k' be the number of chords in the path Q . Then k' is between $a^2 - (2a - 1) = (a - 1)^2$ and $a(a + 1) - 2a = a^2 - a$. In particular,

$$k' \geq (a - 1)^2 \geq (\sqrt{k} - 4)^2 \geq k - 8\sqrt{k}.$$

Let k'' be the number of chords in the path $uwQxx^-$. Then, as $e_G(\{x^-, u\}, V(Q)) \leq 2(2a + 1)$,

$$k'' = k' + e_G(\{x^-, u\}, V(Q)) - 2 \leq a^2 + 3a \leq (a + 2)^2 \leq k.$$

Take $b = k - k''$, so that $0 \leq b \leq 8\sqrt{k}$. Let P' be the subpath of P that starts at x and ends at the $(b + 1)$ -th neighbour u_b of u in $P \setminus x$. Then $uwQxP'u_bu$ is a cycle with exactly k chords. \square

Lemma 6.5. *Let $\ell \gg k \gg 1$, $p \geq 3$, and let K and K' be vertex-disjoint induced copies of $K_{\ell, \ell}$, with no edges between them, in a p -extraction of a graph G . For a unimodal path P that starts in K and is not in the last two layers, let u be a vertex with a neighbour $w \in K' \setminus V(P)$. Then either there is a cycle in G with exactly k chords, or u sends at most $30\sqrt{k}$ edges to P .*

Proof. Let x, y be the ends of P , where $x \in K$, and let x^-, y^- be the neighbours of x, y in P . Take z to be any vertex in K other than x and take v to be any vertex in K' in the opposite side to w . Let Q be a unimodal path with ends z and v from a later layer than P and a different layer than u , and let z^-, v^- be the neighbours of z, v in Q , respectively. By choice of Q , there are no edges between $P \setminus \{x, x^-, y, y^-\}$ and Q (see Observation 3.2).




Figure 7: Lemma 6.5

Suppose that there is no cycle with exactly k chords and that u has more than $30\sqrt{k}$ neighbours in P . By Lemma 6.4, there are at most $8\sqrt{k}$ edges between x^- and Q and at most $8\sqrt{k}$ edges between u and Q . Let Q' be a path in K with ends x and z on either $2a$ or $2a + 1$ vertices, where

$a = \lfloor \sqrt{k} - 13 \rfloor$. Denote by k' the number of chords in the path $R := uvvQzQ'xx^-$, excluding the edge ux^- if it exists. Then

$$k' \geq \#\{\text{chords in } Q'\} \geq (a-1)^2 \geq (\sqrt{k}-15)^2 \geq k - 30\sqrt{k}.$$

As x^-, z^-, v^- , and u are the only vertices in $V(R)$ that may have neighbours in Q' ,

$$\begin{aligned} k' &\leq \#\{\text{chords in } Q'\} + e_G(\{x^-, z^-, v^-, u\}, V(Q')) + e_G(\{x^-, u\}, V(Q)) + e_G(\{x^-, z^-, v^-, v, u, w\}) \\ &\leq (a^2 - a) + 4 \cdot (2a + 1) + 16\sqrt{k} + \binom{5}{2} \leq (a+4)^2 + 16\sqrt{k} \leq (\sqrt{k}-9)^2 + 16\sqrt{k} \\ &= k - 2\sqrt{k} + 81 \leq k, \end{aligned}$$

where the last inequality follows from the assumption that k is large. Take $b = k - k'$, so that $0 \leq b \leq 30\sqrt{k}$. Let P' be the subpath of P that starts at x and ends at the $(b+1)$ -th neighbour u_b of u in $P \setminus \{x, x^-\}$. Then, as there are no edges between $P \setminus \{x, x^-, y, y^-\}$ and K, K' or Q , the cycle $uvvQzQ'xP'u_bu$ has exactly k chords. \square

6.2 Proof of Lemma 6.3

We now prove Lemma 6.3. Roughly speaking, the idea is to find many disjoint induced $K_{\ell, \ell}$'s with no edges between them, and join them via unimodal paths, to obtain a cycle whose length we can control by choosing subpaths of the $K_{\ell, \ell}$'s of appropriate lengths. Variants of the arguments used in this proof will appear in the following section.

Proof of Lemma 6.3. Let K_0, \dots, K_{100} be a collection of vertex-disjoint induced copies of $K_{\ell, \ell}$ in the p -extraction of G such that there are no edges across distinct copies. Denote the bipartition of K_i by $\{U_{i,1}, U_{i,2}\}$. Let $P_{i,j}$ be vertex-disjoint unimodal paths from $U_{i,j}$ to $U_{0,j}$, for $i \in [100]$ and $j \in [2]$. We take all the unimodal paths from different layers that are not the last two. Let $u_{i,j}$ be the first internal vertex on $P_{i,j}$ from K_i , for each $i \in [100]$ and $j \in [2]$. Denote by $u'_{i,j}$ the unique neighbour of $u_{i,j}$ in $V(P_{i,j}) \cap V(K_i)$. That is, the end vertex of $P_{i,j}$ in K_i . A vertex u is said to be *complete* (resp. *anti-complete*) to $U_{i,j}$ if it is adjacent to all (resp. none) of the vertices in $U_{i,j} \setminus \{u'_{i,j}\}$, for $i \in [100]$ and $j \in [2]$.

We first claim that, by possibly shrinking ℓ to $\ell' = \ell/2^{200}$, we may assume that each $u_{i,j}$ is either complete or anti-complete to $U_{s,t}$, for each $i, s \in [100]$ and $j, t \in [2]$. For each $v \in \bigcup_{s=1}^{100} (U_{s,1} \cup U_{s,2})$, write a 0-1 vector $x_v \in \{0, 1\}^{200}$ to encode its adjacency to $u_{i,j}$. That is, the (i, j) -coordinate is 1 if $vu_{i,j} \in E(G)$ and 0 otherwise. Then each $U_{s,t} \setminus \{u_{s,t}\}$ can be partitioned into at most 2^{200} subsets according to the value of x_v . Replacing $U_{s,t}$ by the largest amongst these subsets and adding $u_{s,t}$ suffices for our purpose.

Let $v_{i,j}$ be the first internal vertex on $P_{i,j}$ from K_0 , and let $v'_{i,j}$ be the unique vertex in $V(P_{i,j}) \cap V(K_0)$. By using the same argument possibly shrinking ℓ even further, we may assume that each

lem17.pdf

Figure 8: Part of the cycle $\mathcal{C}(a_1, \dots, a_{100})$

$v_{i,j}$ is either complete or anti-complete to $U_{s,t}$, for $i, s \in [100]$ and $j, t \in [2]$.

Our plan is to make a cycle in the following way. We will choose integers $a_1, \dots, a_{100} \geq 0$, depending on k and the structure we have just found. Starting from $v'_{1,1} \in U_{0,1}$, we take $P_{1,1}$ to reach $u'_{1,1} \in U_{1,1}$, then go through a path of length $2a_1 + 1$ in K_1 to reach $u'_{1,2} \in U_{1,2}$. The journey goes back to K_0 through $P_{1,2}$. Then we move to $v'_{2,1}$ to iterate. After 100 iterations, we close the cycle by moving from $v'_{100,2}$ to $v'_{1,1}$ by an edge.

Let us calculate how many chords exist in such a cycle, which we call $\mathcal{C}(a_1, \dots, a_{100})$ (note that the number of chords depends only on a_1, \dots, a_{100} , and not on the choice of the paths in K_i for $i \in [100]$, by previous assumptions). Firstly, a $(2a_i + 1)$ -edge path in a copy of K_{a_i, a_i} gives $(a_i + 1)^2 - (2a_i + 1) = a_i^2$ chords and the path in K_0 has at most 100^2 chords. Second, each $u_{i,j}$ or $v_{i,j}$ that is complete to $U_{s,t}$ contributes a_s chords (to $U_{s,t} \setminus \{u'_{s,t}\}$), and there are at most 100^2 chords between vertices $u_{i,j}$ or $v_{i,j}$ and the path in K_0 . Third, each $u_{i,j}$ or $v_{i,j}$ adjacent to $u'_{s,t}$, with $(i, j) \neq (s, t)$, contributes one chord. Finally, there are at most $O(\sqrt{k})$ chords between internal vertices of $P_{i,j}$ s (by Lemmas 6.4 and 6.5, because otherwise there is a cycle with exactly k chords and then, we are done). Overall, the cycle $\mathcal{C}(a_1, \dots, a_{100})$ has the following number of chords

$$\sum_{s=1}^{100} a_s^2 + \sum_{s=1}^{100} t_s a_s + O(\sqrt{k}), \quad (1)$$

where t_s is the number of vertices $u_{i,j}$ or $v_{i,j}$ that are complete to $U_{s,1}$ plus the number of vertices $u_{i,j}$ or $v_{i,j}$ that are complete to $U_{s,2}$, and $O(\sqrt{k})$ is a function that only depends on the graph structure that we have found, i.e., the 101 induced $K_{\ell, \ell}$'s and the unimodal paths in between.

In (1), the $O(\sqrt{k})$ term and the terms t_s , with $s \in [100]$, are all fixed parameters, i.e., they do not depend on the choice of a_1, \dots, a_{100} . Also $t_s \leq 400$ for each $s \in [100]$. Thus, one may write (1) as

$$f_G(k) + \sum_{s=1}^{100} (a_s^2 + t_s a_s), \quad (2)$$

where $f_G(k) = O(\sqrt{k})$ is a parameter depending on k and G , but not on a_1, \dots, a_{100} .

Among t_1, \dots, t_{100} , at least 20 values are even, or at least 80 are odd. If the former happens, we may assume t_1, \dots, t_{20} are the even numbers by relabelling the indices. Then, by Lemma 5.1, we

can choose $b_s \geq 200$ for $s \in [20]$, such that

$$\sum_{s=1}^{20} b_s^2 = k - f_G(k) + \sum_{s=1}^{20} \frac{t_s^2}{4}.$$

Indeed, large enough k guarantees the right-hand side is a large enough positive integer that can be expressed as the sum of exactly 20 squares larger than 200. Now let $a_s = b_s - t_s/2$ for $s \in [20]$ and 0 otherwise (notice that $a_2 \geq 200 - t_s/2 \geq 0$). Then the cycle $\mathcal{C}(a_1, a_2, \dots, a_{100})$ has exactly k chords, since (2) becomes

$$f_G(k) + \sum_{s=1}^{100} (a_s^2 + t_s a_s) = f_G(k) + \sum_{s=1}^{20} \left(b_s^2 - \frac{t_s^2}{4} \right) = k.$$

Otherwise, there are at least 80 odd t_s 's, say t_1, \dots, t_{80} . Let r be the unique integer divisible by 4 such that

$$r - 3 \leq k - f_G(k) + \sum_{s=1}^{80} \frac{t_s^2 - 1}{4} \leq r.$$

By our choice of t_1, \dots, t_{80} , $(t_s^2 - 1)/4$ is an integer for each $s \in [80]$. Lemma 5.2 then shows that, since r is a large enough integer divisible by 4, there exist integers $b_s \geq 300$, for $s \in [80]$, such that

$$r = \sum_{s=1}^{80} b_s(b_s + 1).$$

Now take $a_s = b_s - (t_s - 1)/2$ if $s \in [80]$ (so $a_s \geq 300 - (t_s - 1)/2 \geq 0$), and 0 otherwise. Then the number k' of chords in $\mathcal{C}(a_1, a_2, \dots, a_{100})$, using (2), is

$$\begin{aligned} k' &= f_G(k) + \sum_{s=1}^{100} (a_s^2 + t_s a_s) \\ &= f_G(k) + \sum_{s=1}^{80} \left(b_s(b_s + 1) - \frac{t_s^2 - 1}{4} \right) = f_G(k) + r - \sum_{s=1}^{80} \frac{t_s^2 - 1}{4}. \end{aligned}$$

By our choice of r , $k \leq k' \leq k + 3$. Therefore, there is a cycle with exactly k' chords for some $k' \in \{k, k + 1, k + 2, k + 3\}$, as required. \square

7 Exact result

In this section we will prove Theorem 1.1, our main result. To do so, we will prove Lemma 7.7 below, which asserts that if a p -extraction of a graph G contains ℓ pairwise disjoint induced copies of $K_{\ell, \ell}$ with no edges between them, where $p, \ell \gg k \gg 1$, then there is a cycle with exactly k chords. Notice that Theorem 1.1 can be deduced from this lemma, along with Corollary 4.2 and Lemma 6.2,

following the proof of Theorem 6.1 in Section 6 and replacing the call to Lemma 6.3 by a call to Lemma 7.7.

7.1 Overview of the proof

The basic idea for the proof of Lemma 7.7 is similar to that of Lemma 6.3. Given ℓ induced copies K_1, \dots, K_ℓ of $K_{\ell, \ell}$ with no edges between them, we clean them up so that we can build cycles, taking paths $P_i \subseteq K_i$ and connecting them by unimodal paths, in such a way that the number of chords depends only on the lengths of the P_i 's. However, as seen in the previous section, in some cases a parity issue may cause this strategy to fail to give the precise desired number of chords.

To fix this, when cleaning up the K_i 's we take into account, for each vertex u in one of the K_i 's and each triple $T = \{j_1, j_2, j_3\}$ of layers (with $j_1 > j_2 > j_3$), a j_1 -father $f_1(u; T)$ of u , a j_2 -father $f_2(u; T)$ of $f_1(u; T)$, and a j_3 -father $f_3(u; T)$ of $f_2(u; T)$ (recall that in the previous section we considered just one father of one vertex per $K_{\ell, \ell}$).

Lemma 7.2 allows us to assume that there are no edges between $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and K_i 's not containing u . An important step towards the proof of Lemma 7.2 is Lemma 7.1. The latter lemma considers a case where there is a collection of 31 pairwise vertex-disjoint induced $K_{\ell, \ell}$'s with no edges between them, and a collection of 31 vertices that are each complete to each of these $K_{\ell, \ell}$'s, or are each complete to one part of each $K_{\ell, \ell}$ and anti-complete to the other. The proof of this lemma follows the usual scheme of building a cycle and controlling the number of its chords by controlling the number of vertices used from each $K_{\ell, \ell}$. The proof here is, in fact, a bit simpler as instead of connecting the $K_{\ell, \ell}$'s using unimodal paths we can use the 31 vertices.

With the above assumption at hand, namely that there are no edges between the set of parents $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and K_i 's not containing u , we now wish to analyse the interaction between any K_i and the sets $\{f_j(u; T) : u \in K_i\}$, $j = 1, 2, 3$, for any fixed T . This is done in Lemma 7.4. The full statement of the lemma is quite long. In short, it allows us to assume that one of six cases holds, each describing a concrete well-structured graph.

We then turn to the proof of Lemma 7.7, which is the main result of the section. By applying Lemma 7.4 to sufficiently many triples of layers, we can assume that one of the six cases mentioned above holds for enough K_i 's and enough layers. In four of these cases (which we analyse simultaneously), we build the cycles as usual and can reach the desired number of chords without encountering parity issues. In the remaining two cases, we require a few small gadgets to adjust the number of chords in case of a parity issue.

7.2 Removing cross edges

As mentioned above, our first step towards Lemma 7.7 is Lemma 7.2 which allows us to assume that, given a collection of induced $K_{\ell, \ell}$'s with no edges between them, there are no edges between

parents of one copy of $K_{\ell,\ell}$ and any other copy of $K_{\ell,\ell}$. Before proving Lemma 7.2, we prove the following lemma which considers the other extreme, where there are many ‘well-connected’ vertices that are each fully joined to at least one part of each such $K_{\ell,\ell}$.

The proof is similar to other proofs in this paper, where we build a cycle using paths from each $K_{\ell,\ell}$. However, instead of using unimodal paths to connect these paths, we use the well-connected vertices, which simplifies the proof.

Lemma 7.1. *Let $\ell \gg k \gg 1$. In a graph G , suppose that K_1, \dots, K_{31} are pairwise vertex-disjoint induced copies of $K_{\ell,\ell}$ with no edges between them. Denote by $\{U_{i,1}, U_{i,2}\}$ the bipartition of K_i . Let X be set of 31 vertices such that one of the following holds:*

- (a) *every $x \in X$ is complete to K_i , for $i \in [31]$ or*
- (b) *every $x \in X$ is complete to $U_{i,1}$ and anti-competes to $U_{i,2}$, for $i \in [31]$.*

Then there is a cycle in G with exactly k chords.

Proof. Suppose first that (a) is the case. Let $x_1, \dots, x_{20} \in X$ be distinct. Let x be the number of edges in the induced subgraph $G[\{x_1, \dots, x_{20}\}]$. For non-negative integers a_1, \dots, a_{20} , let Q_i be a path in K_i of length $2a_i + 1$. Then each x_i is adjacent to the ends of Q_j for $i, j \in [20]$. Let \mathcal{C} to be the cycle $x_1 Q_1 x_2 \dots x_{20} Q_{20} x_1$.




Figure 9: Part of the cycle \mathcal{C} in Case (a)

We now wish to count the number of chords in \mathcal{C} , which only depends on a_1, \dots, a_{20} and x . First, by counting the edges induced on $V(Q_i)$, the edges from x_i to Q_j , and the edges induced on $\{x_1, \dots, x_{20}\}$, the number of edges with both ends in $V(\mathcal{C})$ is

$$\sum_{i=1}^{20} (a_i + 1)^2 + \sum_{i=1}^{20} 20 \cdot 2(a_i + 1) + x,$$

whereas the length of the cycle is $\sum_{i=1}^{20} (2a_i + 3)$. Subtracting the cycle length from the number of edges induced on $V(\mathcal{C})$, the number of chords equals to

$$\sum_{i=1}^{20} (a_i^2 + 40a_i + 38) + x = \sum_{i=1}^{20} (a_i + 20)^2 + x - c,$$

1em20b.pdf

Figure 10: Part of the cycle \mathcal{C} in Case (b)

where $c = 20 \cdot (400 - 38)$. By using Lemma 5.1 for large enough k , there exist integers $b_1, \dots, b_{20} \geq 20$ such that $\sum_{i=1}^{20} b_i^2 = k - x + c$. Given such b_1, \dots, b_{20} , take $a_i = b_i - 20$ for $i \in [20]$. Then the cycle \mathcal{C} has exactly k chords, as required.

Suppose now that (b) happens. Let $x_1, \dots, x_{21} \in X$ be distinct. Let x be the number of edges in the induced subgraph $G[\{x_1, \dots, x_{21}\}]$. Given integers $a_1, \dots, a_{20} \geq 0$, let Q_i be a path in K_i of length $2a_i$, with both ends in $U_{i,1}$. Let \mathcal{C} be the cycle $x_1 Q_1 x_2 \dots x_{21} Q_{21} x_1$.

Then the number of edges with both ends in the cycle \mathcal{C} is

$$\sum_{i=1}^{21} a_i(a_i + 1) + \sum_{i=1}^{21} 21(a_i + 1) + x,$$

where the length of the cycle is $\sum_{i=1}^{21} (2a_i + 2)$. Thus, the number of chords in \mathcal{C} , obtained by subtracting the number of the \mathcal{C} -edges from the number of induced edges on $V(\mathcal{C})$, is

$$\sum_{i=1}^{21} (a_i^2 + 20a_i + 19) + x = \sum_{i=1}^{21} (a_i + 10)^2 + x - c,$$

where $c = 21 \cdot (100 - 19)$. Again by Lemma 5.1, there exist $b_1, \dots, b_{21} \geq 10$ such that $\sum_{i=1}^{21} b_i^2 = k - x + c$. Given such b_1, \dots, b_{21} , choosing $a_i = b_i - 10$ yields a cycle \mathcal{C} with exactly k chords. \square

The next lemma allows us to assume that there is a collection of many disjoint induced $K_{\ell, \ell}$'s with no edges between them, such that for many triples of layers $T = \{j_1 > j_2 > j_3\}$ and every vertex u in one of these $K_{\ell, \ell}$'s, there are no edges between $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and $K_{\ell, \ell}$'s not containing u (where $f_1(u; T)$ is a j_1 -father of u , $f_2(u; T)$ is a j_2 -father of $f_1(u; T)$, $f_3(u; T)$ is a j_3 -father of $f_2(u; T)$, and $T = \{j_1 > j_2 > j_3\}$).

Roughly speaking, to prove this lemma we choose a collection $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ for each u and T . We then clean up the $K_{\ell, \ell}$'s using Ramsey's theorem to assume that each $f_j(u; T)$ is either complete or anti-complete to each part of each $K_{\ell, \ell}$. Finally, using the previous lemma we may assume that for the vast majority of choices of u , T and $K_{\ell, \ell}$ -copy K , there are no edges between $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and K . The Kövari–Sós–Turán theorem then allows us to find the required structure.

Lemma 7.2. *Let $p, \ell \gg m \gg k \gg 1$ and let G_p be a p -extraction of G . Suppose that, in G_p ,*

K_1, \dots, K_ℓ are pairwise vertex-disjoint induced copies of $K_{\ell, \ell}$ with no edges between them. Then either there is a cycle with exactly k chords, or there exist a collection K'_1, \dots, K'_m of pairwise vertex-disjoint induced $K_{m, m}$'s in G_p and a subset $J \subseteq [p]$ of m indices such that, for every triple $T = \{j_1, j_2, j_3\} \subseteq J$ with $j_1 > j_2 > j_3$, every $i \in [m]$ and every $u \in V(K'_i)$, there exist vertices $f_1(u; T), f_2(u; T), f_3(u; T)$ as follows:

- $f_1(u; T)$ is a j_1 -father of u ,
- $f_2(u; T)$ is a j_2 -father of $f_1(u; T)$,
- $f_3(u; T)$ is a j_3 -father of $f_2(u; T)$,
- there are no edges between $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and $\bigcup_{s \neq i} V(K'_s)$.

Proof. We shall choose a, b, ℓ' with $\ell, p \gg M \gg a \gg b \gg m \gg \ell' \gg k$. For each $u \in K_1 \cup \dots \cup K_\ell$ and a triple $T = \{j_1, j_2, j_3\}$ in $[a]$ (with $j_1 > j_2 > j_3$), choose $f_i(u; T)$, for $i \in [3]$, arbitrarily from those vertices such that $f_1(u; T)$ is a j_1 -father of u and $f_i(u; T)$ is a j_i -father of $f_{i-1}(u; T)$ for $i \in \{2, 3\}$. Let $F(u)$ be the collection of all $f_i(u; T)$ chosen.

Let $\{X_i, Y_i\}$ be the bipartition of K_i . We claim that for $i \in [b]$ we can find subsets $X'_i \subseteq X_i$ and $Y'_i \subseteq Y_i$ of size m so that for all $u \in X'_i \cup Y'_i$ and $j \neq i$, each $v \in F(u)$ is either complete or anti-complete to X'_j and either complete or anti-complete to Y'_j .

This can be done by an analogous technique to the one used in the proof of Lemma 6.3. Namely, for a vertex w and a set Z , let $\mathbf{x}(w, Z) \in \{0, 1\}^Z$ be the 0-1 vector that encodes the adjacency between vertices in Z and w . That is, the corresponding coordinate to $v \in Z$ is 1 if w and v are adjacent and 0 otherwise. Then one can partition X_i (resp. Y_i) according to the values of $\mathbf{x}(w, Z)$ with $w \in X_i$ (resp. Y_i). By choosing the largest subset in the partition, we have subsets $X'_i \subseteq X_i$ and $Y'_i \subseteq Y_i$ such that $|X'_i| \geq 2^{-|Z|}|X_i|$ and $|Y'_i| \geq 2^{-|Z|}|Y_i|$, and each $z \in Z$ is either complete or anti-complete to each $w \in X'_i \cup Y'_i$.

First, choose arbitrary subsets $X_b^1 \subseteq X_b$ and $Y_b^1 \subseteq Y_b$ with $|X_b^1| = |Y_b^1| = M$. Then for each $i < b$, it is possible to choose $X_i^1 \subseteq X_i$ and $Y_i^1 \subseteq Y_i$ of size M such that each vertex $v \in F(u)$, with $u \in X_j^1 \cup Y_j^1$ and $j > i$, is either complete or anti-complete to $X_i^1 \cup Y_i^1$ (indeed, the previous paragraph tells us that there exist such subsets of size at least $\ell \cdot 2^{-a^3 b M} \geq M$).

Next, we repeat in the reverse order. That is, starting with $i = 1$, we choose subsets $X'_i \subseteq X_i^1$ and $Y'_i \subseteq Y_i^1$ of size m such that each vertex $v \in F(u)$, with $u \in X_j^1 \cup Y_j^1$ and $j < i$, is either complete or anti-complete to $X'_i \cup Y'_i$ (this is possible since we are guaranteed such subsets of size at least $M \cdot 2^{-a^3 b m} \geq m$). The sets X'_i and Y'_i satisfy the requirements.

We then collect all ‘fathers’ $v \in F(u)$ for some $u \in X'_i \cup Y'_i$. That is, we set $F := \bigcup_{i=1}^b \bigcup_{u \in X'_i \cup Y'_i} F(u)$. Consider the auxiliary bipartite graph B between F and $[b]$ where $v \in F$ and $j \in [b]$ are adjacent whenever v is complete to X'_j or Y'_j . Colour each edge (v, j) by red, blue, or green if v is complete to X'_j only, Y'_j only, or both $X'_j \cup Y'_j$, respectively. By Lemma 7.1, if there is no cycle with exactly

k chords, then B contains no monochromatic $K_{\ell', \ell'}$. Since $m \gg \ell'$, the bipartite Ramsey Theorem tells us that B contains no copy of $K_{m, m}$. In particular, there are at most $m \binom{b}{m}$ vertices $f \in F$ with $\deg_B(f) \geq m$. Since $a \gg m \binom{b}{m} + m$, we can choose a set J of m indices so that for $j \in J$, all j -fathers $f \in F$ have $\deg_B(f) < m$. For each $u \in \bigcup_{i=1}^b X'_i \cup Y'_i$, let $F'(u) \subseteq F(u)$ be the subset of fathers $f_i(u; T)$ with $i \in \{1, 2, 3\}$ and $T \in \binom{J}{3}$. Note that $|F'(u)| \leq |J|^3 \leq m^3$.

Consider an auxiliary directed graph D on $[b]$ where we join i to j if there is some $u \in X'_i \cup Y'_i$ and $v \in F'(u)$ with v complete to X'_j or Y'_j . As $|\bigcup_{u \in X'_i \cup Y'_i} F'(u)| \leq |X'_i \cup Y'_i| m^3 = 2m^4$ for each i and each $f \in F'(u)$ is adjacent to at most m of the sets $X_j \cup Y_j$, the digraph D has maximum out-degree at most $2m^5 \leq b/(2m+1)$. Therefore, D has an independent set I of size m . Letting K'_1, \dots, K'_m be the complete bipartite graphs induced by $X'_i \cup Y'_i$ for $i \in I$, we get a set of complete bipartite graphs satisfying the lemma. \square

7.3 Analysing the structure of a complete bipartite subgraph and its parents

Our next task is to prove Lemma 7.4, which will be used to analyse the interaction between a complete bipartite subgraph and its parents, ‘grandparents’ and ‘great-grandparents’.

The following simple lemma will be useful in what follows.

Lemma 7.3. *Let G_2 be a bipartite graph on the bipartition $A \cup B$, where $|A| = |B| = m$, and let G_1 be its subgraph. Suppose that $d_{G_1}(a) \geq 1$ for every $a \in A$ and $d_{G_2}(b) \leq d$ for every $b \in B$. Then there exist $A' \subseteq A$ and $B' \subseteq B$, such that $|A'| = |B'| \geq m/4d^2$ and each induced subgraph $G_i[A', B']$, $i = 1, 2$, is a perfect matching.*

Proof. Let $A_0 := \{a \in A : d_{G_2}(a) \leq 2d\}$. Then $e(G_2) \leq d|B|$ implies $|A \setminus A_0| \leq e(G_2)/2d \leq |B|/2 = m/2$ and hence, $|A_0| \geq m/2$. Let $A' \subseteq A_0$ be a maximal subset such that there exists $B' \subseteq B$ such that each $G_i[A', B']$, $i = 1, 2$, induces a perfect matching. We claim that every vertex in $A_0 \setminus A'$ has a G_2 -neighbour in $N_{G_2}(A')$. Indeed, if $a \in A_0 \setminus A'$ has no G_2 -neighbour in $N_{G_2}(A')$ then it has a G_1 -neighbour $b \in B \setminus N_{G_2}(A')$, which makes each $G_i[A' \cup \{a\}, B' \cup \{b\}]$, $i = 1, 2$, a perfect matching. This contradicts the maximality of A' . Thus, $A_0 \subseteq N_{G_2}(N_{G_2}(A'))$, which implies $|A_0| \leq d|N_{G_2}(A')| \leq 2d^2|A'|$. As $|A_0| \geq m/2$, $|B'| = |A'| \geq m/4d^2$, as desired. \square

Lemma 7.4. *For $\ell \gg \omega$, let K be an induced copy of $K_{\ell, \ell}$ with bipartition $X \cup Y$. Suppose that R, S, T are pairwise disjoint sets outside K such that every vertex in K has a neighbour in R , every vertex in R has a neighbour in S , and every vertex in S has a neighbour in T . Let $U := R \cup S \cup T$ and $\ell' := \log \ell / 10000$.*

Then there exist sets X' and Y' of size ℓ' , each of which contained in a different set amongst X and Y , sets $A, A', A'' \subseteq U$, each of which contained in a different set amongst R, S, T , and vertices $a, b \in U$ such that one of the following holds:

(K1) *The vertex a is complete to $X' \cup Y'$.*

- (K2) The vertex a is complete to X' and anti-complete to Y' , and b is complete to Y' and anti-complete to X' .
- (K3) Both the induced bipartite graphs $G[A, X']$ and $G[B, Y']$ are perfect matchings, A is anti-complete to Y' , and B is anti-complete to X' .
- (K4) The induced bipartite graph $G[A, X']$ is a perfect matching and A is complete to Y' .
- (K5) The induced bipartite graph $G[A, X']$ is a perfect matching, A is anti-complete to Y' , A' is anti-complete to X' and either complete or anti-complete to Y' , and every vertex in A has a neighbour in A' .
- (K6) Each component in $G[A, A', A'', Y']$ consists of four vertices u, u', u'', y in A, A', A'', Y' , respectively, such that all pairs in $\{u, u', u'', y\}$, except for possibly uu'' , form edges. Additionally, $A \cup A' \cup A''$ is anti-complete to X' , and b is complete to X' and anti-complete to Y' .

Proof. We begin by proving the following claim.

Claim 7.5. Let $\ell_0 := \ell^{1/7}/10$. There exist $R_1, S_1, T_1 \subseteq U$ of size ℓ_0 , each of which is contained in a different set amongst $R, S, T, Q_1 \subseteq X$ of size ℓ_0 , and a vertex $u_1 \in U$ such that one of the following holds:

- (a) The vertex u_1 is complete to Q_1 .
- (b) The vertex u_1 is complete to R_1 and anti-complete to Q_1 and $G[Q_1, R_1]$ induces a perfect matching.
- (c) Every component in $G[Q_1, R_1, S_1]$ is an induced path qrs with q, r, s in Q_1, R_1, S_1 , respectively.
- (d) Every component in $G[Q_1, R_1, S_1, T_1]$ consists of four vertices q, r, s, t from Q_1, R_1, S_1, T_1 , respectively, such that all pairs in $\{q, r, s, t\}$ are edges, except for possibly rt .

Proof of the claim. We shall use $\ell = (10\ell_0)^7$ implicitly throughout the proof.

If a vertex in U has at least ℓ_0 neighbours in X then we may take u_1 to be this vertex and Q_1 to be a set of ℓ_0 of its neighbours in X , which satisfies (a). In this case, the sets R_1, S_1, T_1 are chosen arbitrarily, as they play no roles in what follows. We may hence assume that every vertex in U sends at most ℓ_0 edges to X .

By Lemma 7.3, there are subsets $Q^{(1)} \subseteq Q$ and $R^{(1)} \subseteq R$ with $|Q^{(1)}| = |R^{(1)}| \geq \ell/4\ell_0^2$ such that $G[Q^{(1)}, R^{(1)}]$ induces a perfect matching. For $r \in R^{(1)}$, denote by $q(r)$ the unique neighbour of r in $Q^{(1)}$; define $r(q)$ similarly for each $q \in Q^{(1)}$.

Suppose that there is a vertex $u \in S \cup T$ with at least $2\ell_0$ neighbours in $R^{(1)}$. As u has at most ℓ_0 neighbours in Q , there are at least ℓ_0 neighbours r of u such that $q(r)$ is not adjacent to u . Let

$R_1 \subseteq R^{(1)}$ be a set of ℓ_0 such vertices r . Then taking $Q_1 := \{q(r) : r \in R_1\}$ and $u_1 := u$ proves (b). Thus, we may assume that every vertex in $S \cup T$ has at most $2\ell_0$ neighbours in $R^{(1)}$.

Let $S^{(1)} \subseteq S$ be a set of size $|R^{(1)}|$ such that each vertex in $R^{(1)}$ has a neighbour in $S^{(1)}$. Let $G_1 = G[R^{(1)}, S^{(1)}]$ and let G_2 be the bipartite graph on $R^{(1)} \cup S^{(1)}$ where $rs, r \in R^{(1)}$ and $s \in S^{(1)}$, is an edge if s has a neighbour in $\{r, q(r)\}$. By Lemma 7.3, there exist subsets $R^{(2)} \subseteq R^{(1)}$ and $S^{(2)} \subseteq S^{(1)}$ with $|S^{(2)}| = |R^{(2)}| \geq |R^{(1)}|/36\ell_0^2 \geq \ell/144\ell_0^4$, such that $G_i[R^{(2)}, S^{(2)}]$ induces a perfect matching, for $i = 1, 2$.

Let $Q^{(2)} := \{q(r) : r \in R^{(2)}\}$ and let $r(s), s(r), q(s), s(q)$ be defined similarly as above for $s \in S^{(2)}$, $r \in R^{(2)}$, and $q \in Q^{(2)}$. That is, $r(s)$ is the unique neighbour of s in $R^{(2)}$; $q(s)$ is the unique neighbour of $r(s)$ in $Q^{(2)}$; etc. Then each component in $G[Q^{(2)}, R^{(2)}, S^{(2)}]$ consists of vertices q, r, s with $q \in Q^{(2)}, r \in R^{(2)}, s \in S^{(2)}$ such that qr and rs are edges. If qs is a non-edge for at least half of the components, then (c) holds. We may thus assume that qs is an edge for at least half of the components, implying that there are subsets $Q^{(3)} \subseteq Q^{(2)}$, $R^{(3)} \subseteq R^{(2)}$ and $S^{(3)} \subseteq S^{(2)}$, with $|Q^{(3)}| = |R^{(3)}| = |S^{(3)}| \geq \ell/288\ell_0^4$, such that $G[Q^{(3)}, R^{(3)}, S^{(3)}]$ induces a K_3 -factor.

Suppose that $u \in T$ has at least $2\ell_0$ neighbours in $S^{(3)}$. Let $R_1 \subseteq S^{(3)}$ be a set of ℓ_0 neighbours such that $q(s)$ is not a neighbour of u . Indeed, such a set exists because u has at most ℓ_0 neighbours in $Q^{(3)}$. Then taking $Q_1 := \{q(s) : s \in R_1\}$ and $u_1 := u$ makes (b) hold. We thus assume that every vertex in T has at most $2\ell_0$ neighbours in $S^{(3)}$.

Now Lemma 7.3 implies that there exist subsets $Q^{(4)}, R^{(4)}, S^{(4)}, T^{(4)}$ of $Q^{(3)}, R^{(3)}, S^{(3)}, T$, respectively, which are of the same size at least $|Q^{(3)}|/100\ell_0^2 \geq \ell/28800\ell_0^6$, such that each component in $G[Q^{(4)}, R^{(4)}, S^{(4)}, T^{(4)}]$ consists of vertices q, r, s, t from Q, R, S, T , respectively, such that qr, qs, rs, ts are edges. If qt is a non-edge for at least half of the components, then (c) holds by taking subsets of $Q^{(4)}, S^{(4)}, T^{(4)}$ that induces 2-edge paths. Otherwise, (d) holds. \square

An analogous claim that produces subsets $Q_2, R_2, S_2, T_2 \subseteq Y$ and vertex u_2 also holds for Y . We may hence fix the vertex sets Q_i, R_i, S_i, T_i and vertices $u_i, i = 1, 2$, that satisfy the claim above and the analogous statement for Y . We then proceed to further refine these subsets.

Claim 7.6. *Let $\ell_1 = \log \ell_0/100$. There exist subset Q'_i, R'_i, S'_i, T'_i of Q_i, R_i, S_i, T_i , respectively, for $i = 1, 2$, which are of the same size ℓ_1 and satisfy one of (the obvious analogues of) (a) to (d), and, additionally, each of u_i, R'_i, S'_i, T'_i is either complete or anti-complete to Q_{3-i} , for $i \in [2]$.*

Proof of the claim. For $q \in Q_1$, let $r(q)$ be the unique neighbour of q in R_1 , assuming one of (b) to (d) holds; otherwise, choose $r(q)$ arbitrarily (say, from R_1 ; it will play no roles). Similarly, let $s(q)$ be the unique neighbour of $r(q)$ in S_1 , if one of (c) and (d) holds, and let $t(q)$ be the unique neighbour of $s(q)$ if (d) holds. For the cases when $s(q)$ or $t(q)$ are not defined, we again choose them arbitrarily as they will play no roles. For $q \in Q_2$, we define $r(q), s(q), t(q)$ analogously with respect to Q_2, R_2, S_2, T_2 .

We then consider an auxiliary edge-coloured complete bipartite graph H on $Q_1 \cup Q_2$ as follows: for $q_1 \in Q_1$ and $q_2 \in Q_2$, colour $q_1 q_2$ by the 0-1 vector of length eight that encodes which of the pairs in $\{q_{3-i}\} \times \{r(q_i), s(q_i), t(q_i), u_i\}$ are edges in G or not, for $i = 1, 2$.

By the classical Kövari–Sós–Turan theorem, there exist subsets $Q'_1 \subseteq Q_1$ and $Q'_2 \subseteq Q_2$ of size ℓ_1 each, such that $H[Q'_1, Q'_2]$ is monochromatic. Taking $R'_i := \{r(q) : q \in Q'_i\}$, $S'_i = \{s(q) : q \in Q'_i\}$, and $T'_i = \{t(q) : q \in Q'_i\}$ proves the claim. \square

For brevity, let us rename Q'_i, R'_i, S'_i, T'_i to Q_i, R_i, S_i, T_i , respectively, for $i = 1, 2$.

A simple (though somewhat tedious) case analysis now shows that one of (K1) to (K6) above holds by using $\ell_1 \geq \log \ell / 10000 = \ell'$.

- **Property (a) holds for $i = 1, 2$.**

We claim that one of (K1) and (K2) holds. Indeed, take $X' := Q_1$ and $Y' := Q_2$. If one of u_1 and u_2 is complete to both Q_1 and Q_2 for some $i \in [2]$, then (K1) hold. Otherwise, taking $a := u_1$ and $b := u_2$ makes (K2) hold.

- **Property (a) holds for neither $i = 1$ nor 2 .**

We claim that one of (K3) and (K4) holds. Indeed, if R_i is anti-complete to Q_{3-i} for both $i = 1$ and 2 , then (K3) holds by taking X', Y', A, B to be Q_1, Q_2, R_1, R_2 , respectively). Otherwise, without loss of generality R_1 is complete to Q_2 , and (K4) holds by taking X', Y', A to be Q_1, Q_2, R_1 .

Without loss of generality, it remains to consider the case where property (a) holds for $i = 2$ but not for $i = 1$. At every step of the following case analysis, we iteratively assume that the previous cases do not hold.

- **The vertex u_2 is complete to Q_1 .** Then (K1) holds.
- **The set R_1 is complete to Q_2 .** Then (K4) holds.
- **Property (b) holds for $i = 1$.**

Then (K5) holds: take X', Y', A, A', b to be $Q_1, Q_2, R_1, \{u_1\}, u_2$.

- **Property (c) holds for $i = 1$.**

Again (K5) holds: take X', Y', A, A', b to be Q_1, Q_2, R_1, S_1, u_2 .

- **Property (d) holds for $i = 1$.**

Here we may assume that R_1, S_1, T_1 are anti-complete to Q_2 , as otherwise (K4) holds. It follows that (K6) holds: take X', Y', A, A', A'', b to be $Q_2, Q_1, R_1, S_1, T_1, u_2$, respectively. \square

7.4 Finding a cycle with exactly k chords

Finally, we prove the following lemma, which finds a cycle with exactly k chords given a large collection of pairwise vertex-disjoint $K_{\ell,\ell}$'s with no edges between them. As mentioned previously, we will use a similar approach to that used in the previous section and earlier in this section, connecting the $K_{\ell,\ell}$'s by unimodal paths and closing a cycle by choosing a path of the right length from each $K_{\ell,\ell}$. The difference here is the much more careful analysis of the interaction between a single $K_{\ell,\ell}$, and a collection of parents, 'grandparents' and 'great-grandparents' of its vertices. This analysis will give rise to three cases (the first corresponding to the first four cases in Lemma 7.4 and the last two each corresponding to one of the last two cases in Lemma 7.4). In the last two cases we will sometimes need to adjust the cycle lengths slightly. For that we build our cycle so as to contain certain small 'gadgets' that will allow us to do so.

Lemma 7.7. *Let $p, \ell \gg k \gg 1$. Suppose that, in a p -extraction of G , there are ℓ pairwise vertex-disjoint induced copies of $K_{\ell,\ell}$ with no edges between them. Then there is a cycle in G with exactly k chords.*

Proof. Suppose that there is no cycle with exactly k chords. Let m satisfy $p, \ell \gg m \gg k$. Then by Lemma 7.2, there exist a collection K_1, \dots, K_{200} of pairwise vertex-disjoint copies of induced $K_{m,m}$'s in the p -extraction of G , a subset $J \subseteq [p]$ of size $200 \cdot 39$ that satisfy: for every triple $T = \{j_1, j_2, j_3\}$ in J with $j_1 > j_2 > j_3$, every $i \in [200]$ and every $u \in K_i$, there exist vertices $f_1(u; T)$, $f_2(u; T)$, and $f_3(u; T)$ such that $f_t(u; T)$ is j_t -father of $f_{t-1}(u; T)$, $t = 1, 2, 3$, where $u = f_0(u; T)$. Moreover, there are no edges between $\{f_1(u; T), f_2(u; T), f_3(u; T)\}$ and $\cup_{s \neq i} V(K_s)$.

Let $S_1, \dots, S_{200} \subseteq J$ be pairwise disjoint subsets of size 39. Fix one S_t and let T_1, \dots, T_{13} be disjoint triples that partition S_t . Now repeatedly apply Lemma 7.4 thirteen times to each K_i : at the beginning, let $K = K_i$ with the bipartition $X_i^{(0)} \cup Y_i^{(0)}$. At the j -th iteration, we apply Lemma 7.4 with $X = X_i^{(j-1)}$, $Y = Y_i^{(j-1)}$, $R = \{f_1(u; T_j) : u \in X_i^{(j-1)} \cup Y_i^{(j-1)}\}$, $S = \{f_2(u; T_j) : u \in X_i^{(j-1)} \cup Y_i^{(j-1)}\}$ and $T = \{f_3(u; T_j) : u \in X_i^{(j-1)} \cup Y_i^{(j-1)}\}$. Then let $X_i^{(j)}$ and $Y_i^{(j)}$ be the resulting subsets $X' \subseteq X$ and $Y' \subseteq Y$, respectively.

During the 13 iterations, we fall into one of the three cases, which we denote by (1), (2) and (3): (1) analyses the case when one of (K1), (K2), (K3) and (K4) occurs at least twice; the remaining cases when (K5) or (K6) occur at least twice are described in (2) and (3), respectively. For brevity, write $X_i = X_i^{(13)}$ and $Y_i = Y_i^{(13)}$; X_i and Y_i are the shrunken sets at the end of the 13 iterations, each of which has size at least $(\log^{(26)} m)/10^6$.⁴

- (1) There are vertices $u_{i,1}$ and $u_{i,2}$ in two distinct layers with indices in S_i such that $u_{i,1}$ and $u_{i,2}$ are anti-complete to $X_s \cup Y_s$ for $s \in [200] \setminus \{i\}$ and one of the following holds. For each case, we also identify $u'_{i,1}$ and $u'_{i,2}$ that will be useful for further analysis.

- $2 \times (\text{K1})$: $u_{i,1}$ and $u_{i,2}$ are complete to $X_i \cup Y_i$; choose $u'_{i,1} \in X_i$ and $u'_{i,2} \in Y_i$.

⁴Here $\log^{(\ell)} x$ denotes ℓ compositions of logarithm, e.g., $\log^{(2)} x = \log \log x$.

- $2 \times (\text{K2})$: $u_{i,1}$ is complete to X_i and anti-complete to Y_i and $u_{i,2}$ is complete to Y_i and anti-complete to X_i ; choose $u'_{i,1} \in X_i$ and $u'_{i,2} \in Y_i$.
- $2 \times (\text{K3})$: each of $u_{i,1}$ and $u_{i,2}$ has exactly one neighbour in $X_i \cup Y_i$, which we denote $u'_{i,1} \in X_i$ and $u'_{i,2} \in Y_i$, respectively.
- $2 \times (\text{K4})$: $u_{i,1}$ and $u_{i,2}$ are complete to Y_i , and each have exactly one neighbour in X_i , which we denote $u'_{i,1}$ and $u'_{i,2}$, respectively, such that $u'_{i,1} \neq u'_{i,2}$.

(2) There are vertices $u_{i,1}, \dots, u_{i,4}$ from four distinct layers in S_i , such that:

- $u_{i,1}, \dots, u_{i,4}$ are anti-complete to $X_j \cup Y_j$, for $j \in \{0, \dots, 200\} \setminus \{i\}$.
- $u_{i,1}$ and $u_{i,3}$ are anti-complete to Y_i , and each of them has a unique neighbour in X_i , denoted by $u'_{i,1}$ and $u'_{i,3}$, respectively, such that $u'_{i,1} \neq u'_{i,3}$.
- $u_{i,2}$ is complete to Y_i and anti-complete to X_i ; let $u'_{i,2} \in Y_i$.
- $u_{i,4}$ is anti-complete to X_i , either complete or anti-complete to Y_i , and is adjacent to $u_{i,3}$.

(3) There are vertices $u_{i,1}, \dots, u_{i,5}$ from five distinct layers in S_i , such that:

- $u_{i,1}, \dots, u_{i,5}$ are anti-complete to $X_j \cup Y_j$ for $j \in \{0, \dots, 200\} \setminus \{i\}$.
- both $u_{i,1}$ and $u_{i,3}$ have exactly one neighbour in X_i , denoted $u'_{i,1}$ and $u'_{i,3}$, respectively, that are distinct, and are anti-complete to Y_i .
- $u_{i,2}$ is complete to Y_i and anti-complete to X_i ; let $u'_{i,2} \in Y_i$.
- $u_{i,4}$ is a neighbour of $u_{i,3}$ and $u'_{i,3}$ and is anti-complete to $(X_i \cup Y_i) \setminus \{u'_{i,3}\}$.
- $u_{i,5}$ is a neighbour of $u'_{i,3}$ and $u_{i,4}$ and is anti-complete to $(X_i \cup Y_i) \setminus \{u'_{i,3}\}$.

To see this, let T_1, \dots, T_{13} be disjoint sets of size 3 in S_i . Apply Lemma 7.4 thirteen times: at the i -th time apply it with the current $X_i^{(j)}$ and $Y_i^{(j)}$, and with the sets $\{f_1(u; T_i) : u \in Z\}$, $\{f_2(u; T_i) : u \in Z\}$ and $\{f_3(u; T_i) : u \in Z\}$, where $Z = X_i^{(j)} \cup Y_i^{(j)}$. If one of (K1), (K2), (K3) and (K4) occurs at least twice, then (1) holds. Indeed, say (K1) holds twice, then we can take $u_{i,1}$ and $u_{i,2}$ to be the vertices a corresponding to some two iterations in which (K1) held. If (K5) occurs four times then (2) holds, and if (K6) occurs five times then (3) holds. Here taking multiple occurrences is essential to guarantee that all $u_{i,1}, \dots, u_{i,5}$ lie in distinct layers. It then follows that one of the following occurs: (1) holds for 20 values of $i \in [200]$; (2) holds for 100 values of $i \in [200]$; or (3) holds for 80 values of $i \in [200]$.

Case (1) holds for at least 20 values of $i \in [200]$.

By relabelling the indices of the 200 copies of $K_{m,m}$ if necessary, we may assume that (1) holds for $i \in [20]$. To construct a cycle with k chords, we need one more copy of $K_{m,m}$, so we shall use K_{21} too. For simplicity, let $K_0 = K_{21}$ be the copy of $K_{m,m}$ on $X_0 \cup Y_0$, where $X_0 = X_{21}$ and $Y_0 = Y_{21}$.

We want to take a set of distinct vertices $w_{i,j}$, $i \in [20]$ and $j = 1, 2$, where each $w_{i,j}$ is at the same layer as $u_{i,j}$ and has a neighbour $w'_{i,j}$ in K_0 which are all distinct too. Here the index $j = 1, 2$

indicates $w'_{i,1} \in X_0$ and $w'_{i,2} \in Y_0$, as was done for $u'_{i,j}$'s. Indeed, this is possible since one can greedily take $w'_{i,1}$ and $w'_{i,2}$ disjoint from the previous choices in X_0 and Y_0 , respectively, and let $w_{i,j} = f_1(w'_{i,j}; T)$ where T is the triple in S_i such that $u_{i,j} = f_1(u'_{i,j}; T)$. Let $P_{i,j}$ be a unimodal path between $u_{i,j}$ and $w_{i,j}$, taken in the same layers as the two end vertices so that they are in different layers from the K_i 's. Note that there are no edges between $w_{i,j}$ and $X_i \cup Y_i$ for $i \in [20]$.

Given integers $\alpha_1, \dots, \alpha_{20}$, we define a cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{20})$ as follows. Let Q_i be the path from $w'_{i,1}$ to $w_{i+1,1}$ for $i \in [20]$ (addition of indices taken modulo 20), obtained by concatenating the paths $w'_{i,1}w_{i,1}$, $P_{i,1}$, $u_{i,1}u'_{i,1}$, Q_i^* , $u'_{i,2}u_{i,2}$, $P_{i,2}$, $w_{i,2}w'_{i,2}w'_{i+1,1}$, where Q_i^* is a path in $G[X_i, Y_i]$ with ends $u'_{i,1}$ and $u'_{i,2}$ of length $2\alpha_i + 1$. Now let $\mathcal{C}(\alpha_1, \dots, \alpha_{20})$ be the cycle obtained by concatenating Q_1, \dots, Q_{20} .




Figure 11: The path Q_i in Case (1)

While the cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{20})$ is not uniquely defined, its number of chords depends only on $\alpha_1, \dots, \alpha_{20}$. To evaluate the number of chords in this cycle, Q_i^* contributes $(\alpha_i + 1)^2 - (2\alpha_i + 1) = \alpha_i^2$ chords; the number of chords with one end in $\{u_{i,1}, u_{i,2}\}$ and one end in $X_i \cup Y_i$ is one of the four values $2(2\alpha_i + 1)$, $2\alpha_i$, 0 , and $2(\alpha_i + 1)$, depending on the four cases described in (1); there are at most 120^2 chords with both ends in $\bigcup_{i \in [20], j \in [2]} \{w_{i,j}, w'_{i,j}, u_{i,j}\}$; and we may assume that the number of chords with ends in $\bigcup_{i \in [20], j \in [2]} V(P_{i,j})$ is $O(\sqrt{k})$ by Lemmas 6.4 and 6.5. Note that there are no chords between distinct sets $X_i \cup Y_i$, between a set $X_i \cup Y_i$ and $\{u_{j,1}, u_{j,2}, w_{j,1}, w_{j,2}\}$ for distinct $i, j \in [20]$, or between $\{u_{i,1}, u_{i,2}\}$ and $\{w_{j,1}, w'_{j,1}, w_{j,2}, w'_{j,2}\}$. In total, the number of chords is

$$f_G(k) + \sum_{i \in [20]} (\alpha_i^2 + 2\sigma_i \alpha_i) = f_G(k) + \sum_{i \in [20]} ((\alpha_i + \sigma_i)^2 - \sigma_i^2) = h_G(k) + \sum_{i \in [20]} (\alpha_i + \sigma_i)^2.$$

where $\sigma_i \in \{0, 1, 2\}$ for $i \in [20]$, and $f_G(k)$ and $h_G(k)$ are functions that do not depend on $\alpha_1, \dots, \alpha_{20}$ and satisfy $f_G(k), h_G(k) = O(\sqrt{k})$. By Lemma 5.1, there is a choice of integers $\beta_1, \dots, \beta_{20} \geq 2$ such that $\sum_{i \in [20]} \beta_i^2 = k - h_G(k)$. A cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{20})$ with $\alpha_i = \beta_i - \sigma_i$ has k chords, as desired.

Case (2) holds for at least 100 values of $i \in [200]$.

Again by possible relabelling, we may assume that (2) holds for $i \in [100]$. We also need an extra copy, denoted by K_0 , taken from K_i , $i > 100$, on the bipartition $X_0 \cup Y_0$. Let $w_{i,j}$, with $i \in [100]$ and $j \in [4]$, satisfy the following: $w_{i,j}$ is at the same layer as $u_{i,j}$; it is anti-complete to $X_i \cup Y_i$ for $i \in [100]$; $w_{i,j}$ has a neighbour $w'_{i,j}$ such that $w'_{i,j} \in X_0$ if $j \in \{1, 3\}$ and otherwise $w'_{i,j} \in Y_0$; and the vertices $w_{i,j}, w'_{i,j}$, with $j \in [4]$ and $i \in [100]$, are all distinct. Indeed, such choices of $w_{i,j}$ and $w'_{i,j}$

1em23b.pdf

Figure 12: The path Q_i in Case (2)

are possible since one can again take $w_{i,j} = f_1(w'_{i,j}; T)$ with the triple T such that $u_{i,j} = f_1(u'_{i,j}; T)$ while maintaining all $w'_{i,j}$ being distinct.

Let $P_{i,j}$ be a unimodal path with ends $u_{i,j}$ and $w_{i,j}$ in the same layer as the end vertices, for each $i \in [100]$ and $j \in [4]$. Given positive integers $\alpha_1, \dots, \alpha_{100}$, we define a cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ as follows. Let Q_i be the path from $w'_{i,1}$ to $w'_{i+1,1}$ (index addition taken modulo 100), obtained by concatenating the paths $w'_{i,1}w_{i,1}$, $P_{i,1}$, $u_{i,1}u'_{i,1}$, Q_i^* , $u'_{i,2}u_{i,2}$, $P_{i,2}$, $w_{i,2}w'_{i,2}w'_{i,3}w_{i,3}$, $P_{i,3}$, $u_{i,3}u_{i,4}$, $P_{i,4}$, $w_{i,4}w'_{i,4}w'_{i+1,1}$, where Q_i^* is a path in $G[X_i \setminus \{u'_{i,3}\}, Y_i]$ with ends $u'_{i,1}$ and $u'_{i,2}$, and of length $2\alpha_i + 1$. Let $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ be the cycle obtained by concatenating Q_1, \dots, Q_{100} .

As above, the cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ is not uniquely defined, but the number of its chords depends only on $\alpha_1, \dots, \alpha_{100}$. Let us evaluate the number of chords precisely as follows: Q_i^* contributes α_i^2 chords; the number of chords between $\{u_{i,1}, \dots, u_{i,4}\}$ and $X_i \cup Y_i$ is either α_i or $2\alpha_i + 1$ depending on how $u_{i,4}$ connects to Y_i (either complete or anti-complete); there are at most 1200^2 chords with ends in $\bigcup_{i \in [100], j \in [4]} \{w_{i,j}, w'_{i,j}, u_{i,j}\}$; and we may assume that the number of chords in $\bigcup_{i \in [100], j \in [4]} V(P_{i,j})$ is $O(\sqrt{k})$ by Lemmas 6.4 and 6.5. In total, the number of chords in the cycle is

$$f_G(k) + \sum_{i \in [100]} (\alpha_i^2 + \sigma_i \alpha_i),$$

where $\sigma_i \in \{1, 2\}$ and $f_G(k) = O(\sqrt{k})$. Here σ_i 's and $f_G(k)$ do not depend on α_i 's.

Suppose that $\sigma_i = 2$ for at least 20 values of $i \in [100]$; let I be a set of 20 such values of i . Set $\alpha_i = 1$ for $i \in [100] \setminus I$. Each α_i , $i \in I$, is not yet determined. Then the number of chords in a cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ reduces to

$$h_G(k) + \sum_{i \in I} (\alpha_i + 1)^2,$$

for some $h_G(k) = O(\sqrt{k})$ that does not depend on α_i 's with $i \in I$. By Lemma 5.1, there exist integers β_i , for $i \in I$, such that $\beta_i \geq 2$ and $\sum_{i \in I} \beta_i^2 = k - h_G(k)$. Set $\alpha_i = \beta_i - 1$. The cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ then has exactly k chords.

It remains to consider the case where $\sigma_i = 1$ for at least 80 values of $i \in [100]$. Let I be a set of 80 values of i such that $\sigma_i = 1$. Set $\alpha_i = 1$ for $i \in [100] \setminus I$. Then the number of chords in the cycle $\mathcal{C}(\alpha_1, \dots, \alpha_{100})$ is

$$h_G(k) + \sum_{i \in I} (\alpha_i^2 + \alpha_i),$$

where $h_G(k) = O(\sqrt{k})$ is independent of the α_i 's. Let $\tau \in \{0, 1, 2, 3\}$ be the remainder of $k - h_G(k)$

Figure 13: The path Q_i in Case (3)

modulo 4. Let $\mathcal{C}_\tau(\alpha_1, \dots, \alpha_{100})$ be a cycle obtained by replacing an inner vertex of $V(Q_i^*) \cap X_i$ by $u'_{i,3}$, for τ values of i . One can readily check that the number of chords in $\mathcal{C}_\tau(\alpha_1, \dots, \alpha_{100})$ is $h_G(k) + \tau + \sum_{i \in I} (\alpha_i^2 + \alpha_i)m$ as we gain the additional chord $u_{i,3}u'_{i,3}$ for τ values of i . Now, since $k - h_G(k) - \tau$ is divisible by 4, Lemma 5.2 implies that there exist integers α_i , $i \in I$, such that $\alpha_i \geq 1$ and $\sum_{i \in I} (\alpha_i^2 + \alpha_i) = k - h_G(k) - \tau$. In particular, there is a cycle with exactly k chords.

Case (3) holds for at least 80 values of $i \in [200]$

Without loss of generality, (3) holds for $i \in [80]$ and let $K_0 = K_{81}$ be another copy of $K_{m,m}$ on $X_0 \cup Y_0$. Analogously to the previous cases, choose $w_{i,j}$, $i \in [80]$ and $j \in [5]$, that satisfy the following conditions: $w_{i,j}$ is in the same layer as $u_{i,j}$; it is anti-complete to $X_i \cup Y_i$ for $i \in [80]$; $w_{i,j}$ has a neighbour $w'_{i,j}$ such that $w'_{i,j} \in X_0$ for $j \in \{1, 3\}$ and $w'_{i,j} \in Y_0$ for $j \in \{2, 4, 5\}$; and the vertices $w_{i,j}, w'_{i,j}$, with $i \in [80]$ and $j \in [5]$, are all distinct. Again, such choices for $w_{i,j}$ and $w'_{i,j}$ exist since one can set $w_{i,j} = f_1(w'_{i,j}, T)$ with the triple T such that $u_{i,j} = f_1(u'_{i,j}; T)$, while maintaining all $w'_{i,j}$ being distinct.

One can find a path $P_{i,j}$ between $u_{i,j}$ and $w_{i,j}$ in the same layer as the two ends, for each $i \in [80]$ and $j \in [5]$. In fact, we do not make use of $w_{i,4}, w'_{i,4}$ and $P_{i,4}$, but we took them to preserve the correspondence between indices.

Given positive integers $\alpha_1, \dots, \alpha_{80}$, we choose a cycle $\mathcal{C}^0(\alpha_1, \dots, \alpha_{80})$ as follows. Let Q_i be the path from $w'_{i,1}$ to $w'_{i+1,1}$ (addition modulo 80), obtained by concatenating the paths $w'_{i,1}w_{i,1}, P_{i,1}, u_{i,1}u'_{i,1}, Q_i^*, u'_{i,2}u_{i,2}, P_{i,2}, w_{i,2}w'_{i,2}w'_{i,3}w_{i,3}, P_{i,3}, u_{i,3}u_{i,4}u_{i,5}, P_{i,5}, w_{i,5}w'_{i,5}w'_{i+1,1}$; where Q_i^* is a path in $G[X_i \setminus \{u_{i,3}\}, Y_i]$ with ends $u'_{i,1}$ and $u'_{i,2}$ and of length $2\alpha_i + 1$. Let $\mathcal{C}^0(\alpha_1, \dots, \alpha_{80})$ be the cycle obtained by concatenating Q_1, \dots, Q_{80} . For $\tau = 1, 2, 3$, let $\mathcal{C}^\tau(\alpha_1, \dots, \alpha_{80})$ be the cycle obtained from $\mathcal{C}^0(\alpha_1, \dots, \alpha_{80})$ by replacing one of the internal vertices of Q_i^* in X_i by $u'_{i,3}$, for $i \in [\tau]$.

The cycle $\mathcal{C}^\sigma(\alpha_1, \dots, \alpha_{80})$ for each $\sigma = 0, 1, 2, 3$ is not necessarily unique, but all the choices have the same number of chords. To verify, let us evaluate the number of chords in $\mathcal{C}^0(\alpha_1, \dots, \alpha_{80})$ first, as follows: Q_i^* contributes α_i^2 chords; there are exactly α_i chords between $\{u_{i,1}, \dots, u_{i,5}\}$ and $X_i \cup Y_i$, all of which are incident to $u_{i,2}$; there are at most 1200^2 chords with ends in $\bigcup_{i \in [80], j \in [5]} \{w_{i,j}, w'_{i,j}, u_{i,j}\}$; and we may assume that the number of chords in $\bigcup_{i \in [80], j \in [5]} V(P_{i,j})$ is $O(\sqrt{k})$, by Lemmas 6.4 and 6.5. The cycle $\mathcal{C}^\tau(\alpha_1, \dots, \alpha_{80})$ has precisely 3τ more chords than $\mathcal{C}^0(\alpha_1, \dots, \alpha_{80})$, as $u'_{i,3}$ gives three extra chords to $u_{i,3}, u_{i,4}$, and $u_{i,5}$ (see Figure 13). The total number of chords in $\mathcal{C}^\tau(\alpha_1, \dots, \alpha_{80})$ is hence

$$f_G(k) + \sum_{i \in [80]} (\alpha_i^2 + \alpha_i) + 3\tau,$$

where $f_G(k) = O(\sqrt{k})$. Now choose $\tau \in \{0, 1, 2, 3\}$ be such that 3τ equals $k - f_G(k)$ modulo 4. By Lemma 5.2, there exist positive integers $\alpha_1, \dots, \alpha_{80}$, such that $\sum_{i \in [80]} \alpha_i(\alpha_i + 1) = k - f_G(k) - 3\tau$. The cycle $\mathcal{C}^\tau(\alpha_1, \dots, \alpha_{80})$ has exactly k chords. \square

8 Conclusion

We showed that the family of graphs with no cycle with exactly k chords is χ -bounded, for every integer k which is either sufficiently large or of form $k = \ell(\ell - 2)$, where $\ell \geq 3$ is an integer. This was already known to hold for $k \in \{1, 2, 3\}$ (see [1, 18]). An obvious follow-up problem, which is a conjecture due to Aboulker and Bousquet [1], would be to extend this to all $k \geq 1$.

Conjecture 8.1 (Aboulker–Bousquet [1]). *For every $k \geq 1$ there is a function f_k such that, if G is a graph with no cycle with exactly k chords, then $\chi(G) \leq f_k(\omega(G))$.*

It would also be interesting to get a better understanding of whether graphs with small clique number and large chromatic number contain cycles with k chords that have some sort of further structure. Whereas our proof essentially just produces a cycle with k chords, there are conjectures that it should be possible to find more. A k -fan is defined as a path P with one additional vertex v added that has exactly k neighbours on P . It is easy to see that k -fans contain a cycle with $k - 2$ chords. Thus the following would be a strengthening of the results in this paper and of Conjecture 8.1.

Conjecture 8.2 (Davies [5]). *For every $k \geq 1$ there is a function f_k such that, if G is a graph with k -fan, then $\chi(G) \leq f_k(\omega(G))$.*

Currently, this conjecture is only known to hold for $k = 1$ (where is it equivalent to the “ $k = 1$ ” case of Conjecture 8.1).

References

- [1] P. Aboulker and N. Bousquet, *Excluding cycles with a fixed number of chords*, Discr. App. Math. **180** (2015), 11–24.
- [2] B. Bollobás and A. Thomason, *Highly linked graphs*, Combinatorica **16** (1996), 313–320.
- [3] N. Bousquet and S. Thomassé, *On the chromatic number of wheel-free graphs with no large bipartite graphs*, 2015.
- [4] J. Chalpion, L. Esperet, Z. Li, and P. Ossona de Mendez, *Restricted frame graphs and a conjecture of Scott*, Electron. J. Combin. **23** (2016), P1.30.
- [5] J. Davies, *Triangle-free graphs with large chromatic number and no induced wheel*, arXiv:2104.05907 (2021).
- [6] P. Erdős, *Graph theory and probability*, Canad. J. Math. **11** (1959), 34–38.

- [7] A. Gyárfás, *On Ramsey covering-numbers*, Infinite and Finite Sets **2** (1975), 801–816.
- [8] ———, *Problems from the world surrounding perfect graphs*, Applicationes Mathematicae **19** (1987), 413–441.
- [9] H. A. Kierstead and S. G. Penrice, *Radius two trees specify χ -bounded classes*, J. Graph Theory **18** (1994), no. 2, 119–129.
- [10] H. A. Kierstead and Y. Zhu, *Radius three trees in graphs with large chromatic number*, SIAM J. Discr. Math. **17** (2004), 571–581.
- [11] D. Kühn and D. Osthus, *Induced subdivisions in $K_{s,s}$ -free graphs of large average degree*, Combinatorica **24** (2004), 287–304.
- [12] J. Mycielski, *Sur le coloriage des graphes*, Colloq. Math, vol. 3, 1955, pp. 161–162.
- [13] A. Pawlik, J. Kozik, T. Krawczyk, M. Lasoń, P. Micek, W. T. Trotter, and B. Walczak, *Triangle-free intersection graphs of line segments with large chromatic number*, J. Combin. Theory Ser. B **105** (2014), 6–10.
- [14] A. D. Scott, *Induced trees in graphs of large chromatic number*, J. Graph Theory **24** (1997), 297–311.
- [15] D. P. Sumner, *Subtrees of a graph and chromatic number*, pp. 557–576, John Wiley & Sons, New York, 1981.
- [16] R. Thomas and P. Wollan, *An improved linear edge bound for graph linkages*, European J. Combin. **26** (2005), 309–324.
- [17] N. Trotignon, *Perfect graphs: a survey*, arXiv:1301.5149 (2013).
- [18] N. Trotignon and K. Vušković, *A structure theorem for graphs with no cycle with a unique chord and its consequences*, J. Graph Theory **63** (2010), 31–67.