

χ -boundedness of graphs
with no cycle with k chords

Shoham Letzter
University College London

Joint work with Joonkyung Lee and Alexey Pokrovskiy

CanADAM
May 2021


Chromatic number vs clique number

1/10

$\omega(G)$ = size of largest clique in G .

$\chi(G)$ = min number of colours in a proper colouring of $V(G)$.

adjacent vertices have different colours



Observation: $\chi(G) \geq \omega(G)$ for every graph G .

The opposite is far from true. E.g.: there exist Δ -free graphs with arbitrarily large χ .

Chudnovsky-Robertson-Seymour-Thomas '06: $\chi(H) = \omega(H)$ for every induced subgraph of G iff neither G nor G^c has induced odd cycle of length ≥ 5 .

Gyárfás '87: A class of graphs \mathcal{G} is χ -bounded if there is a function f s.t. $\forall G \in \mathcal{G}$

$\chi(G) \leq f(\omega(G))$.

closed under taking induced subgraphs.

Gyárfás '87: A class \mathcal{G} is χ -bounded if there is f s.t. $\forall G \in \mathcal{G} \chi(G) \leq f(\omega(G))$.

$\text{Forb}(H) = \{ \text{graphs with no induced copy of } H \}$.

- If H contains a cycle then $\text{Forb}(H)$ is not χ -bounded.

Erdős '59: \exists graphs with arbitrarily large χ and girth.

\curvearrowright length of shortest cycle

- $\text{Forb}(S_k)$ is χ -bounded $\forall k$.

\curvearrowright star on k edges

Let $G \in \text{Forb}(S_k)$, $u \in V(G)$.


$N(u)$ has no clique of size $\omega(G)$ or independent set of size k .

$|N(u)| \leq r(R_{\omega(G)}, K_k) \forall u \Rightarrow \chi(G) \leq r(R_{\omega(G)}, K_k) + 1$.

\curvearrowright Ramsey number

- Gyárfás '87: $\text{Forb}(P_k)$ is χ -bounded $\forall k$.

\curvearrowright path on k vertices



Forbidding a tree

3/10

- $\text{Forb}(H)$
- is χ -bounded if H is a path or star and
 - isn't χ -bounded if H contains a cycle.


Conjecture (Gyárfás '87): $\text{Forb}(H)$ is χ -bounded iff H is a forest.
($\Leftrightarrow \text{Forb}(T)$ is χ -bounded \forall tree T .)

- Kierstead-Penrice '94: $\text{Forb}(T)$ is χ -bounded \forall tree T of radius ≤ 2 .
- Kierstead-Zhu '06: $\text{Forb}(T)$ is χ -bounded \forall tree T of radius ≤ 3 .

$\text{Forb}^*(H) = \{ \text{graphs with no induced copy of a subdivision of } H. \}$

- Scott '97: $\text{Forb}^*(T)$ is χ -bounded \forall tree T .


Question: For which H is $\text{Forb}^*(H)$ χ -bounded?



$\mathcal{C}_k = \{\text{graphs with no cycle with exactly } k \text{ chords}\}.$

Conjecture (Aboulker-Bousquet '15): \mathcal{C}_k is χ -bounded $\forall k$.

- Trotignon-Vušković '10: \mathcal{C}_1 is χ -bounded.
- Aboulker-Bousquet '15: \mathcal{C}_2 and \mathcal{C}_3 are χ -bounded.



a cycle with
3 chords

Theorem (Lee-L.-Pokrovskiy '21+):

There exists k_0 s.t. \mathcal{C}_k is χ -bounded $\forall k \geq k_0$.

Finding complete bipartite graphs

5/10

Need to show: if $\chi(G) \gg \omega(G), k$ then G has a cycle with exactly k chords.


Theorem: $\forall k, l \exists f$ s.t. if $\chi(G) \geq f(\omega(G))$ then G contains an induced $K_{l,l}$ or a cycle with k chords.

(Uses Bousquet-Thomasse '15: $\text{Forb}(\Delta, K_{l,l}, W_k)$ is χ -bounded $\forall k, l$.)
 \uparrow k -wheel

A cycle C of length $2a$ in a $K_{l,l}$ has $a^2 - 2a$ chords.

#edges induced by $V(C)$ \downarrow edges in C

\Rightarrow If $k = a^2 - 2a$ for some integer a then \exists cycle with k chords.



Corollary: $\forall k, l \exists f$ s.t. if $\chi(G) \geq f(\omega(G))$ then G contains l disjoint induced $K_{l,l}$'s with no edges between them or a cycle with k chords.

Unimodal paths


6/10

Fix $u \in V(G)$. $N_i(u) = \{v : \text{dist}(u, v) = i\}$.

$$\chi(G) \leq \max_i \{\chi(N_i(u) \cup N_{i+1}(u))\} \leq \max_i \{\chi(N_i(u)) + \chi(N_{i+1}(u))\}$$

$\Rightarrow \exists i$ s.t. $\chi(N_i(u)) \geq \frac{1}{2} \cdot \chi(G)$. Write $G_1 = G[N_1(u)]$.

Observation: $\forall x, y \in V(G_1) \exists$ induced path from x to y whose interior is in $V(G) \setminus V(G_1)$.
Such a path is called unimodal.




Many unimodal paths

7/10

Repeat the process to obtain $G = G_0 \supseteq G_1 \supseteq \dots \supseteq G_p$ s.t.:

- $\chi(G_p) \geq 2^{-p} \chi(G)$
- $\forall x, y \in V(G_p) \exists$ unimodal path from x to y with interior in $V(G_i) \setminus V(G_p)$.



Combining complete bipartite graphs

8/10

K_1, \dots, K_{100} : disjoint K_{a_i} 's in G_p with no edges between them.


Take P_i a path of length $2a_i$ in K_i .

Join by unimodal paths from distinct G_j 's.

May assume: \forall part X of K_i and u here u is adjacent to all or none of X .

Lemma: $\exists O(\sqrt{k})$ chords here

$$\Rightarrow \# \text{ chords in depicted cycle} = \sum_{i=1}^{100} (a_i^2 + \underbrace{\sigma_i}_{O(1)} a_i) + \underbrace{O(\sqrt{k})}_{\text{independent of } a_i\text{'s}}.$$



$\forall a_1, \dots, a_{100} \exists$ cycle with the following number of chords: $\sum_{i=1}^{100} (a_i^2 + \sigma_i a_i) + O(\sqrt{k})$.

Lemma. $\forall c$, large k : $\exists a_1, \dots, a_{20} \geq c$ s.t. $k = a_1^2 + \dots + a_{20}^2$.

Lemma. $\forall c$, large k s.t. $4|k$: $\exists a_1, \dots, a_{80} \geq c$ s.t. $k = (a_1^2 + a_1) + \dots + (a_{80}^2 + a_{80})$.

Apply one of above lemmas (depending on parity of σ_i) to find a cycle with k' chords, where $k' \in \{k, k-1, k-2, k-3\}$.

To get exactly k chords we work much harder to find gadgets to offset parity.


Conjecture (Aboulker-Bousquet '15): C_k is χ -bounded $\forall k$.
 graphs with no cycle with k chords

Known for $k=1,2,3$; we proved for large k and $k=a^2-2a, a \in \mathbb{N}$.

A wheel is an induced cycle C plus a vertex adjacent to ≥ 3 vertices of C .

Conjecture (Trotignon '13): {wheel-free graphs} is χ -bounded.

Aboulker-Bousquet '15: {wheel & triangle-free} is χ -bounded.



Conjecture (Gyárfás '87): $\text{Forb}(T)$ is χ -bounded \forall tree T .

Question: when is $\text{Forb}^*(H)$ χ -bounded?

Thank you for your attention!